

5

UNIQUE ASSOCIATED KAPOSI'S SARCOMA VIRUS SEQUENCES AND  
USES THEREOF

10 The invention disclosed herein was made with  
Government support under a co-operative agreement  
CCU210852 from the Centers for Disease Control and  
Prevention, of the Department of Health and Human  
Services. Accordingly, the U.S. Government has  
15 certain rights in this invention.

*hcl* → This application is a continuation-in-part application  
of U.S. Serial No. 08/420,235, filed on April 11, 1995  
which is a continuation-in-part application of U.S.  
20 Serial No. 08/343,101, filed on November 21, 1994, a  
continuation-in-part application of U.S. Serial No.  
08/292,365, filed on August 18, 1994, which is hereby  
incorporated by reference.

25 Throughout this application, various publications may  
be referenced by Arabic numerals in brackets. Full  
citations for these publications may be found at the  
end of each Experimental Details Section. The  
disclosures of the publications cited herein are in  
30 their entirety hereby incorporated by reference into  
this application to more fully describe the state of  
the art to which this invention pertains.

BACKGROUND OF THE INVENTION

35 Kaposi's sarcoma (KS) is the most common neoplasm  
occurring in persons with acquired immunodeficiency  
syndrome (AIDS). Approximately 15-20% of AIDS  
patients develop this neoplasm which rarely occurs in  
immunocompetent individuals [13, 14]. Epidemiologic  
40 evidence suggests that AIDS-associated KS (AIDS-KS)

006230 6472050

has an infectious etiology. Gay and bisexual AIDS patients are approximately twenty times more likely than hemophiliac AIDS patients to develop KS, and KS may be associated with specific sexual practices among  
5 gay men with AIDS [6, 15, 55, 83]. KS is uncommon among adult AIDS patients infected through heterosexual or parenteral HIV transmission, or among  
10 pediatric AIDS patients infected through vertical HIV transmission [77]. Agents previously suspected of causing KS include cytomegalovirus, hepatitis B virus, human papillomavirus, Epstein-Barr virus, human herpesvirus 6, human immunodeficiency virus (HIV), and Mycoplasma penetrans [18, 23, 85, 91, 92]. Non-  
15 infectious environmental agents, such as nitrite inhalants, also have been proposed to play a role in KS tumorigenesis [33]. Extensive investigations, however, have not demonstrated an etiologic association between any of these agents and AIDS-KS [37, 44, 46, 90].

005250 02F40960

SUMMARY OF THE INVENTION

5 This invention provides an isolated DNA molecule which is at least 30 nucleotides in length and which uniquely defines a herpesvirus associated with Kaposi's sarcoma. This invention provides an isolated herpesvirus associated with Kaposi's sarcoma.

10 This invention provides a method of vaccinating a subject for KS, prophylaxis diagnosing or treating a subject with KS and detecting expression of a DNA virus associated with Kaposi's sarcoma in a cell.

RECEIVED

**BRIEF DESCRIPTION OF THE FIGURES****Figure 1:**

5 Agarose gel electrophoresis of RDA products from  
AIDS-KS tissue and uninvolved tissue. RDA was  
performed on DNA extracted from KS skin tissue  
and uninvolved normal skin tissue obtained at  
autopsy from a homosexual man with AIDS-KS. Lane  
10 1 shows the initial PCR amplified genomic  
representation of the AIDS-KS DNA after Bam HI  
digestion. Lanes 2-4 show that subsequent cycles  
of ligation, amplification, hybridization and  
digestion of the RDA products resulted in  
15 amplification of discrete bands at 380, 450, 540  
and 680 bp. RDA of the extracted AIDS-KS DNA  
performed against itself resulted in a single  
band at 540 bp (lane 5). Bands at 380 bp and 680  
bp correspond to KS330Bam and KS627Bam  
20 respectively after removal of 28 bp priming  
sequences. Bands at 450 and 540 bp hybridized  
nonspecifically to both KS and non-KS human DNA.  
Lane M is a molecular weight marker.

**25 Figures 2A-2B:**

Hybridization of <sup>32</sup>P-labelled KS330Bam (Figure 2A)  
and KS627Bam (Figure 2B) sequences to a  
representative panel of 19 DNA samples extracted  
from KS lesions and digested with Bam HI.  
30 KS330Bam hybridized to 11 of the 19 and KS627Bam  
hybridized to 12 of the 19 DNA samples from AIDS-  
KS lesions. Two additional cases (lanes 12 and  
13) were shown to have faint bands with both  
KS330Bam and KS627Bam probes after longer  
35 exposure. One negative specimen (lane 3) did not  
have microscopically detectable KS in the tissue

specimen. Seven of 8 additional KS DNA samples also hybridized to both sequences.

**Figures 3A-3F:**

5 Nucleotide sequences of the DNA herpesvirus associated with KS (KSHV).

**Figures 4A-4B:**

10 PCR amplification of a representative set of KS-derived DNA samples using KS330<sub>234</sub> primers. Figure 4A shows the agarose gel of the amplification products from 19 KS DNA samples (lanes 1-19) and Figure 4B shows specific hybridization of the PCR products to a <sup>32</sup>P end-labelled 25 bp internal oligonucleotide (Figure 3B) after transfer of the gel to a nitrocellulose filter. Negative samples in lanes 3 and 15 respectively lacked microscopically detectable KS in the sample or did not amplify the constitutive p53 exon 6, suggesting that these samples were negative for technical reasons. An additional 8 AIDS-KS samples were amplified and all were positive for KS330<sub>234</sub>. Lane 20 is a negative control and Lane M is a molecular weight marker.

25

**Figure 5:**

30 Southern blot hybridization of KS330Bam and KS627Bam to AIDS-KS genomic DNA extracted from three subjects (lanes 1, 2, and 3) and digested with PvuII. Based on sequence information (Figure 3A), restricted sites for Pvu II occur between bp 12361-12362 of the KSHV sequence (Figure 3A, SEQ ID NO: 1), at bp 134 in KS330Bam (Figure 3B, SEQ ID NO: 2) and bp 414 in KS627Bam (Figure 3C, SEQ ID NO: 3). KS330Bam and KS627Bam failed to hybridize to the same fragments in the digests indicating that the two sequences are

35

5 separated from each other by one or more intervening Bam HI restriction fragments. Digestion with Pvu II and hybridization to KS330Bam resulted in two distinct banding patterns (lanes 1 and 2 vs. lane 3) suggesting variation between KS samples.

**Figure 6:**

10 Comparison of amino acid homologies between EBV ORF BDLF1, HSVSA ORF 26 and a 918 bp reading frame of the Kaposi's sarcoma agent which includes KS330Bam. Amino acid identity is denoted by reverse lettering. In HSVSA, ORF 26 encodes a minor capsid VP23 which is a late gene product.

15

000290" 6274960  
 SUB  
 B1



**Figure 7:**

20 Subculture of Raji cells co-cultivated with BCBL-1 cells treated with TPA for 2 days. PCR shows that Raji cells are positive for KSHV sequences and indicate that the agent is a transmissible virus.

**Figure 8:**

25 A schematic diagram of the orientation of KSHV open reading frames identified on the KS5 20,710 bp DNA fragment. Homologs to each open reading frame from a corresponding region of the herpesvirus saimiri (HSVSA) genome are present in an identical orientation, except for the region corresponding to the ORF 28 of HSVSA (middle schematic section). The shading for each open reading frame corresponds to the approximate % amino acid identity for the KSHV ORF compared to this homolog in HSVSA. Noteworthy homologs that are present in this section of DNA include homologs to thymidine kinase (ORF21), gH

30

35

glycoprotein (ORF22), major capsid protein (ORF25) and the VP23 protein (ORF26) which contains the original KS330Bam sequence derived by representational difference analysis.

5

**Figure 9:**

The ~200 kD antigen band appearing on a Western blot of KS patient sera against BCBL1 lysate (B1) and Raji lysate (RA). M is molecular weight marker. The antigen is a doublet between ca. 210 kD and 240 kD.

10

**Figure 10:**

5 control patient sera without KS (A1N, A2N, A3N, A4N and A5N). B1=BCBL1 lysate, RA=Raji lysate. The 220 kD band is absent from the Western blots using patient sera without KS.

15

**Figure 11:**

In this figure, 0.5 ml aliquots of the gradient have been fractionated (fractions 1-62) with the 30% gradient fraction being at fraction No. 1 and the 10% gradient fraction being at fraction No. 62. Each fraction has been dot hybridized to a nitrocellulose membrane and then a <sup>32</sup>P-labeled KSHV DNA fragment, KS631Bam has been hybridized to the membrane using standard techniques. The figure shows that the major solubilized fraction of the KSHV genome bands (i.e. is isolated) in fractions 42 through 48 of the gradient with a high concentration of the genome being present in fraction 44. A second band of solubilized KSHV DNA occurs in fractions 26 through 32.

20

25

30

35

**Figure 12:**

Location, feature, and relative homologies of KS5 open reading frames compared to translation

products of herpesvirus saimiri (HSV), equine herpesvirus 2 (EHV2) and Epstein-Barr virus (EBV).

5 Figure 13:

Indirect immunofluorescence end-point and geometric mean titers (GMT) in AIDS-KS and AIDS control sera against BHL-6 and P3H3 prior to and after adsorption with P3H3.

10

Figure 14:

Genetic map of KS5, a 20.7 kb lambda phage clone insert derived from a human genomic library prepared from an AIDS-KS lesion. Seventeen partial and complete open reading frames (ORFs) are identified with arrows denoting reading frame orientations. Comparable regions of the Epstein-Barr virus (EBV) and herpesvirus saimiri (HVS) genomes are shown for comparison. Levels of amino acid similarity between KSHV ORFs are indicated by shading of EBV and HVS ORFs (black, over 70% similarity; dark gray, 55-70% similarity; light gray, 40-54% similarity; white, no detectable homology). Domains of conserved herpesvirus sequence blocks and locations of restriction endonuclease sites used in subcloning are shown beneath the KSHV map (B, Bam HI site; N, Not I site). The small Bam HI fragment (black) in the VP23 gene homolog corresponds to the KS330Bam fragment generated by representational difference analysis which was used to identify the KS5 lambda phage clone.

15

20

25

30

Figures 15A-15B:

Phylogenetic trees of KSHV based on comparison of aligned amino acid sequences between herpesviruses for the MCP gene and for a

35

000000000000000000000000

SUB  
P2



concatenated nine-gene set. The comparison of  
MCP sequences (Figure 15A) was obtained by the  
neighbor-joining method and is shown in unrooted  
form with branch lengths proportional to  
divergence (mean number of substitution events  
per site) between the nodes bounding each branch.  
Comparable results were obtained by maximum  
parsimony analysis. The number of times out of  
100 bootstrap samplings the division indicated by  
each internal branch was obtained are shown next  
to each branch; bootstrap values below 75 are not  
shown. Figure 15B is a phylogenetic tree of  
gammaherpesvirus sequences based on a nine-gene  
set CS1 (see text) and demonstrates that KSHV is  
most closely related to the gamma-2 herpesvirus  
sublineage, genus Rhadinovirus. The CS1 amino  
acid sequence was used to infer a tree by the  
Protml maximum likelihood method; comparable  
results, not shown were obtained with the  
neighbor-joining and maximum parsimony methods.  
The bootstrap value for the central branch is  
marked. On the basis of the MCP analysis, the  
root must lie between EBV and the other three  
species. Abbreviations for virus species used in  
the sequence comparisons are 1)  
Alphaherpesvirinae: HSV1 and HSV2, herpes  
simplex virus types 1 and 2; EHV1, equine  
herpesvirus 1; PRV, pseudorabies virus; and VZV,  
varicella-zoster virus, 2) Betaherpesvirinae:  
HCMV, human cytomegalovirus; HHV6 and HHV7, human  
herpesviruses 6 and 7, and 3) Gammaherpesvirinae:  
HVS, herpesvirus saimiri; EHV2, equine  
herpesvirus 2; EBV, Epstein-Barr virus; and  
Kaposi's sarcoma-associated herpesvirus.

**Figures 16A-16B:**

CHEF gel electrophoresis of BCBL-1 DNA hybridized to KS631Bam (Figure 16A) and EBV terminal repeat (Figure 16B). KS631Bam hybridizes to a band at 270 kb as well as to a diffuse band at the origin. The EBV termini sequence hybridizes to a 150-160 kb band consistent with the linear form of the genome. Both KS631Bam (dark arrow) and an EBV terminal sequence hybridize to high molecular weight bands immediately below the origin indicating possible concatemeric or circular DNA. The high molecular weight KS631Bam hybridizing band reproduces poorly but is visible on the original autoradiographs.

**Figure 17:**

Induction of KSHV and EBV replication in BCBL-1 with increasing concentrations of TPA. Each determination was made in triplicate after 48 h of TPA incubation and hybridization was standardized to the amount of cellular DNA by hybridization to beta-actin. The figure shows the mean and range of relative increase in hybridizing genome for EBV and KSHV induced by TPA compared to uninduced BCBL-1. TPA at 20 ng/ml induced an eight-fold increase in EBV genome (upper line) at 48 h compared to only a 1.4 fold increase in KSHV genome (lower line). Despite the lower level of KSHV induction, increased replication of KSHV genome after induction with TPA concentrations over 10 ng/ml was reproducibly detected.

**Figures 18A-18C:**

*In situ* hybridization with an ORF26 oligomer to BCBL-1, Raji and RCC-1 cells. Hybridization occurred to nuclei of KSHV infected BCBL-1

5 (Figure 18A), but not to uninfected Raji cells (Figure 18B). RCC-1, a Raji cell line derived by cultivation of Raji with BCBL-1 in communicating chambers separated by a 0.45  $\mu$  filter, shows rare cells with positive hybridization to the KSHV ORF26 probe (Figure 18C).

Figures 19A-19D:

10 Representative example of IFA staining of BHL-6 with AIDS-KS patient sera and control sera from HIV-infected patients without KS. Both AIDS-KS (Figure 19A) and control (Figure 19B) sera show homogeneous staining of BHL-6 at 1:50 dilution. After adsorption with paraformaldehyde-fixed P3H3 to remove cross-reacting antibodies directed against lymphocyte and EBV antigens, antibodies from AIDS-KS sera localize to BHL-6 nuclei (Figure 19C). P3H3 adsorption of control sera eliminates immunofluorescent staining of BHL-6 (Figure 19D).

Figures 20A-20B:

25 Longitudinal PCR examination for KSHV DNA of paired PBMC samples from AIDS-KS patients (A) and homosexual/bisexual AIDS patients without KS (B). Time 0 is the date of KS onset for cases or other AIDS-defining illness for controls. All samples were randomized and examined blindly. Overall, 7 of the KS patients were KSHV positive at both examination dates (solid bars) and 5 converted from a negative to positive PBMC sample (forward striped bars) immediately prior to or after KS onset. Two previously positive KS patients were negative after KS diagnosis (reverse striped bars) and the remaining KS patients were negative at both timepoints (open bars). Two homosexual/bisexual control PBMC samples without

095040550  
 SUB  
 33

KS converted from negative to positive and one control patient reverted from PCR positive to negative for KSHV DNA.

5 **Figure 21:**  
Sample collection characteristics for AIDS-KS patients, gay/bisexual AIDS patients and hemophilic AIDS patients.

10 **Figure 22:**  
PCR analysis of KS330<sub>233</sub> in DNA samples from patients with Kaposi's sarcoma and tumor controls.

15 **Figure 23:**  
Characteristics of the study population of patients with KS and without KS.

20 **Figure 24:**  
Prevalence of antibody to KSHV p40 in HIV-1 positive patients with and without KS.

25 **Figure 25:**  
Comparison of KS patients with and without antibody to KSHV p40.

30 **Figure 26:**  
Prevalence of antibody detectable by indirect immunofluorescence to KSHV antigens in chemically induced BCBL-1 cells in HIV-1 positive patients with and without KS.

35 **Figures 27A-27B:**  
Specific recognition of KSHV polypeptides in chemically treated BCBL-1 cells. Figure 27A shows reactivity of untreated BCBL-1 and B95-8 cells with RM, a reference human antibody to EBV.

RM recognizes the EBV polypeptides EBNA1 and p21 in the BCBL-1 cells. Figure 27B shows reactivity of untreated and chemically treated cells with serum 01-03 from a patient with KS. Cells were treated with TPA and n-butyrate for 48 hrs. For description of the cell lines see Materials and Methods. The immunoblots were prepared from 10% SDS polyacrylamide gels.

Figures 28A-28D:

Detection of KSHV p40 by sera from patients with KS. Extracts were prepared from BCBL-1 cells (containing KSHV and EBV) and Clone HH514-16 cells (containing only EBV) that were uninduced or treated for 48 hrs with chemical inducing agents, n-butyrate, TPA, or a combination of the two chemicals. Immunoblots prepared from 12% SDS polyacrylamide gels were reacted with a 1:200 dilution of serum from HIV-1 positive patients. Figure 28A shows serum 01-06 from a patient with KS. Figure 28B shows serum 01-07 from a patient without KS. Figure 28C shows serum 04-01 from a patient with KS. Figure 28D shows serum 01-03 from a patient with KS.

Figures 29A-29F:

Detection of KSHV lytic cycle antigens by indirect immunofluorescence. BCBL-1 cells were untreated (Figures 29A, 29C, and 29E) or treated with n-butyrate (Figures 29B, 29D, and 29F) for 48 hrs. Indirect immunofluorescence with a 1:10 dilution of serum from two patients with KS, 04-18 (Figures 29A, and 29B) and 04-38 (Figures 29E, and 29F) and a serum, 04-37 (Figures 29C, and 29D), from a patient without KS.

DETAILED DESCRIPTION OF THE INVENTIONDefinitions

5 The following standard abbreviations are used throughout the specification to indicate specific nucleotides:

10 C=cytosine                    A=adenosine  
T=thymidine                   G=guanosine

15 The term "nucleic acids", as used herein, refers to either DNA or RNA. "Nucleic acid sequence" or "polynucleotide sequence" refers to a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. It includes both self-replicating plasmids, infectious polymers of DNA or RNA and nonfunctional DNA or RNA.

20 By a nucleic acid sequence "homologous to" or "complementary to", it is meant a nucleic acid that selectively hybridizes, duplexes or binds to viral DNA sequences encoding proteins or portions thereof when the DNA sequences encoding the viral protein are present in a human genomic or cDNA library. A DNA  
25 sequence which is homologous to a target sequence can include sequences which are shorter or longer than the target sequence so long as they meet the functional test set forth. Hybridization conditions are  
30 specified along with the source of the CDNA library.

Typically, the hybridization is done in a Southern blot protocol using a 0.2XSSC, 0.1% SDS, 65°C wash. The term "SSC" refers to a citrate-saline solution of  
35 0.15 M sodium chloride and 20 Mm sodium citrate. Solutions are often expressed as multiples or fractions of this concentration. For example, 6XSSC

0022915272056

refers to a solution having a sodium chloride and sodium citrate concentration of 6 times this amount or 0.9 M sodium chloride and 120 mM sodium citrate. 0.2XSSC refers to a solution 0.2 times the SSC concentration or 0.03 M sodium chloride and 4 mM sodium citrate.

The phrase "nucleic acid molecule encoding" refers to a nucleic acid molecule which directs the expression of a specific protein or peptide. The nucleic acid sequences include both the DNA strand sequence that is transcribed into RNA and the RNA sequence that is translated into protein. The nucleic acid molecule include both the full length nucleic acid sequences as well as non-full length sequences derived from the full length protein. It being further understood that the sequence includes the degenerate codons of the native sequence or sequences which may be introduced to provide codon preference in a specific host cell.

The phrase "expression cassette", refers to nucleotide sequences which are capable of affecting expression of a structural gene in hosts compatible with such sequences. Such cassettes include at least promoters and optionally, transcription termination signals. Additional factors necessary or helpful in effecting expression may also be used as described herein.

The term "operably linked" as used herein refers to linkage of a promoter upstream from a DNA sequence such that the promoter mediates transcription of the DNA sequence.

The term "vector", refers to viral expression systems, autonomous self-replicating circular DNA (plasmids), and includes both expression and nonexpression plasmids. Where a recombinant microorganism or cell

culture is described as hosting an "expression vector," this includes both extrachromosomal circular DNA and DNA that has been incorporated into the host chromosome(s). Where a vector is being maintained by a host cell, the vector may either be stably replicated by the cells during mitosis as an autonomous structure, or is incorporated within the host's genome.

The term "plasmid" refers to an autonomous circular DNA molecule capable of replication in a cell, and includes both the expression and nonexpression types. Where a recombinant microorganism or cell culture is described as hosting an "expression plasmid", this includes latent viral DNA integrated into the host chromosome(s). Where a plasmid is being maintained by a host cell, the plasmid is either being stably replicated by the cells during mitosis as an autonomous structure or is incorporated within the host's genome.

The phrase "recombinant protein" or "recombinantly produced protein" refers to a peptide or protein produced using non-native cells that do not have an endogenous copy of DNA able to express the protein. The cells produce the protein because they have been genetically altered by the introduction of the appropriate nucleic acid sequence. The recombinant protein will not be found in association with proteins and other subcellular components normally associated with the cells producing the protein.

The following terms are used to describe the sequence relationships between two or more nucleic acid molecules or polynucleotides: "reference sequence", "comparison window", "sequence identity", "percentage of sequence identity", and "substantial identity". A



"reference sequence" is a defined sequence used as a basis for a sequence comparison; a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length cDNA or gene sequence given in a sequence listing or may comprise a complete cDNA or gene sequence.

Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith and Waterman (1981) *Adv. Appl. Math.* 2:482, by the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. (USA)* 85:2444, or by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, WI).

As applied to polypeptides, the terms "substantial identity" or "substantial sequence identity" mean that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap which share at least 90 percent sequence identity, preferably at least 95 percent sequence identity, more preferably at least 99 percent sequence identity or more.

"Percentage amino acid identity" or "percentage amino acid sequence identity" refers to a comparison of the amino acids of two polypeptides which, when optimally aligned, have approximately the designated percentage of the same amino acids. For example, "95% amino acid identity" refers to a comparison of the amino acids of two polypeptides which when optimally aligned have 95% amino acid identity. Preferably, residue positions which are not identical differ by conservative amino

acid substitutions. For example, the substitution of amino acids having similar chemical properties such as charge or polarity are not likely to effect the properties of a protein. Examples include glutamine for asparagine or glutamic acid for aspartic acid.

The phrase "substantially purified" or "isolated" when referring to a herpesvirus peptide or protein, means a chemical composition which is essentially free of other cellular components. It is preferably in a homogeneous state although it can be in either a dry or aqueous solution. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high performance liquid chromatography. A protein which is the predominant species present in a preparation is substantially purified. Generally, a substantially purified or isolated protein will comprise more than 80% of all macromolecular species present in the preparation. Preferably, the protein is purified to represent greater than 90% of all macromolecular species present. More preferably the protein is purified to greater than 95%, and most preferably the protein is purified to essential homogeneity, wherein other macromolecular species are not detected by conventional techniques.

The phrase "specifically binds to an antibody" or "specifically immunoreactive with", when referring to a protein or peptide, refers to a binding reaction which is determinative of the presence of the herpesvirus of the invention in the presence of a heterogeneous population of proteins and other biologics including viruses other than the herpesvirus. Thus, under designated immunoassay conditions, the specified antibodies bind to the herpesvirus antigens and do not bind in a significant

amount to other antigens present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, antibodies raised to the human herpesvirus immunogen described herein can be selected to obtain antibodies specifically immunoreactive with the herpesvirus proteins and not with other proteins. These antibodies recognize proteins homologous to the human herpesvirus protein. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See Harlow and Lane [32] for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity.

"Biological sample" as used herein refers to any sample obtained from a living organism or from an organism that has died. Examples of biological samples include body fluids and tissue specimens.

I. Kaposi's Sarcoma (KS) - Associated Herpesvirus.

This invention provides an isolated DNA molecule which is at least 30 nucleotides in length and which uniquely defines a herpesvirus associated with Kaposi's sarcoma.

In one embodiment the isolated DNA molecule comprises at least a portion of the nucleic acid sequence as shown in Figure 3A (SEQ ID NO: 1). In another embodiment the isolated DNA molecule is a 330 base pair (bp) sequence. In another embodiment the isolated DNA molecule is a 12-50 bp sequence. In

another embodiment the isolated DNA molecule is a 30-37 bp sequence.

In another embodiment the isolated DNA molecule is genomic DNA. In another embodiment the isolated DNA molecule is cDNA. In another embodiment a RNA is derived from the isolated nucleic acid molecule or is capable of hybridizing with the isolated DNA molecule. As used herein "genomic" means both coding and non-coding regions of the isolated nucleic acid molecule.

Further, the DNA molecule above may be associated with lymphoproliferative diseases including, but not limited to: Hodgkin's disease, non-Hodgkin's lymphoma, lymphatic leukemia, lymphosarcoma, splenomegaly, reticular cell sarcoma, Sezary's syndrome, mycosis fungoides, central nervous system lymphoma, AIDS related central nervous system lymphoma, post-transplant lymphoproliferative disorders, and Burkitt's lymphoma. A lymphoproliferative disorder is characterized as being the uncontrolled clonal or polyclonal expansion of lymphocytes involving lymph nodes, lymphoid tissue and other organs.

This invention provides an isolated nucleic acid molecule encoding an ORF20 (SEQ ID NOs: 22 and 23), ORF21 (SEQ ID NOs:14 and 15), ORF22 (SEQ ID NOs:16 and 17), ORF23 (SEQ ID NOs:18 and 19), ORF24 (SEQ ID NOs: 20 and 21), ORF25 (SEQ ID NOs: 2 and 3), ORF26 (SEQ ID NOs:24 and 25), ORF27 (SEQ ID NOs:26 and 27), ORF28 (SEQ ID NOs:28 and 29), ORF29A (SEQ ID NOs:30 and 31), ORF29B (SEQ ID NOs:4 and 5), ORF30 (SEQ ID NOs:6 and 7), ORF31 (SEQ ID NOs:8 and 9), ORF32 (SEQ ID NOs:32 and 33), ORF33 (SEQ ID NOs: 10 and 11), ORF34 (SEQ ID NOs: 34 and 35), or ORF35 (SEQ ID NOs:12 AND 13).

This invention provides an isolated polypeptide encoded by ORF20 (SEQ ID NOS: 22 and 23), ORF21 (SEQ ID NOS:14 and 15), ORF22 (SEQ ID NOS:16 and 17), ORF23 (SEQ ID NOS:18 and 19), ORF24 (SEQ ID NOS: 20 and 21),  
 5 ORF25 (SEQ ID NOS: 2 and 3), ORF26 (SEQ ID NOS:24 and 25), ORF27 (SEQ ID NOS:26 and 27), ORF28 (SEQ ID NOS:28 and 29), ORF29A (SEQ ID NOS:30 and 31), ORF29B (SEQ ID NOS:4 and 5), ORF30 (SEQ ID NOS:6 and 7), ORF31 (SEQ ID NOS:8 and 9), ORF32 (SEQ ID NOS:32 and  
 10 33), ORF33 (SEQ ID NOS: 10 and 11), ORF34 (SEQ ID NOS: 34 and 35), or ORF35 (SEQ ID NOS:12 AND 13).

For Example, TK is encoded by ORF 21; glycoprotein H (gH) by ORF 22; major capsid protein (MCP) by ORF 25;  
 15 virion polypeptide (VP23) by ORF 26; and minor capsid protein by ORF 27.

This invention provides for a replicable vector comprising the isolated DNA molecule of the DNA virus.  
 20 The vector includes, but is not limited to: a plasmid, cosmid,  $\lambda$  phage or yeast artificial chromosome (YAC) which contains at least a portion of the isolated nucleic acid molecule.

As an example to obtain these vectors, insert and vector DNA can both be exposed to a restriction enzyme to create complementary ends on both molecules which base pair with each other and are then ligated  
 25 together with DNA ligase. Alternatively, linkers can be ligated to the insert DNA which correspond to a restriction site in the vector DNA, which is then digested with the restriction enzyme which cuts at that site. Other means are also available and known  
 30 to an ordinary skilled practitioner.

35

Regulatory elements required for expression include promoter or enhancer sequences to bind RNA polymerase

and transcription initiation sequences for ribosome binding. For example, a bacterial expression vector includes a promoter such as the lac promoter and for transcription initiation the Shine-Dalgarno sequence and the start codon AUG. Similarly, a eukaryotic expression vector includes a heterologous or homologous promoter for RNA polymerase II, a downstream polyadenylation signal, the start codon AUG, and a termination codon for detachment of the ribosome. Such vectors may be obtained commercially or assembled from the sequences described by methods well-known in the art, for example the methods described above for constructing vectors in general.

This invention provides a host cell containing the above vector. The host cell may contain the isolated DNA molecule artificially introduced into the host cell. The host cell may be a eukaryotic or bacterial cell (such as E.coli), yeast cells, fungal cells, insect cells and animal cells. Suitable animal cells include, but are not limited to Vero cells, HeLa cells, Cos cells, CV1 cells and various primary mammalian cells.

This invention provides an isolated herpesvirus associated with Kaposi's sarcoma. In one embodiment the herpesvirus comprises at least a portion of a nucleotide sequence as shown in Figures 3A (SEQ ID NO: 1).

In one embodiment the herpesvirus may be a DNA virus. In another embodiment the herpesvirus may be a Herpesviridae. In another embodiment the herpesvirus may be a gammaherpesvirinae. The classification of the herpesvirus may vary based on the phenotypic or molecular characteristics which are known to those skilled in the art.

This invention provides an isolated DNA virus wherein the viral DNA is about 270 kb in size, wherein the viral DNA encodes a thymidine kinase, and wherein the viral DNA is capable of selectively hybridizing to a nucleic acid probe selected from the group consisting of SEQ ID NOs: 38-40.

The KS-associated human herpesvirus of the invention is associated with KS and is involved in the etiology of the disease. The taxonomic classification of the virus has not yet been made and will be based on phenotypic or molecular characteristics known to those of skill in the art. However, the novel KS-associated virus is a DNA virus that appears to be related to the Herpesviridae family and the gammaherpesvirinae subfamily, on the basis of nucleic acid homology.

A. Sequence identity of the viral DNA and its proteins.

The human herpesvirus of the invention is not limited to the virus having the specific DNA sequences described herein. The KS-associated human herpesvirus DNA shows substantial sequence identity, as defined above, to the viral DNA sequences described herein. DNA from the human herpesvirus typically selectively hybridizes to one or more of the following three nucleic acid probes:

Probe 1 (SEQ ID NO:38)

AGCCGAAAGG ATTCCACCAT TGTGCTCGAA TCCAACGGAT TTGACCCCGT  
GTTCCCATG GTCGTGCCGC AGCAACTGGG GCACGCTATT CTGCAGCAGC  
TGTTGGTGTA CCACATCTAC TCCAAAATAT CGGCCGGGGC CCCGGATGAT  
GTAATATGG CGGAACTTGA TCTATATACC ACCAATGTGT CATTATGGG  
GCGCACATAT CGTCTGGACG TAGACAACAC GGA

Probe 2 (SEQ ID NO:39):

GAAATTACCC ACGAGATCGC TTCCTGCAC ACCGCACTTG GCTACTCATC  
 AGTCATCGCC CCGGCCACG TGGCCGCCAT AACTACAGAC ATGGGAGTAC  
 ATTGTCAGGA CCTCTTTATG ATTTTCCCAG GGGACGCGTA TCAGGACCGC  
 5 CAGCTGCATG ACTATATCAA AATGAAAGCG GGCCTGCAAA CCGGCTCACC  
 GGGAAACAGA ATGGATCACG TGGGATACAC TGCTGGGGTT CCTCGCTGCG  
 AGAACCTGCC CGGTTTGAGT CATGGTCAGC TGGCAACCTG CGAGATAATT  
 CCCACGCCGG TCACATCTGA CGTTGCCT

10

Probe 3 (SEQ ID NO: 40):

AACACGTCAT GTGCAGGAGT GACATTGTGC CGCGGAGAAA CTCAGACCGC  
 ATCCCGTAAC CACTGTGAGT GGGAAAATCT GCTGGCTATG TTTTCTGTGA  
 TTATCTATGC CTTAGATCAC AACTGTCACC CG

15

Hybridization of a viral DNA to the nucleic acid  
 probes listed above is determined by using standard  
 nucleic acid hybridization techniques as described  
 herein. In particular, PCR amplification of a viral  
 20 genome can be carried out using the following three  
 sets of PCR primers:

- 1) ~~AGCCGAAAGGATTCCACCAT;  
TCCGTGTTGTCTACGTCCAG (SEQ ID NO: 41)~~
- 2) ~~GAAATTACCCACGAGATCGC;  
AGGCAACGTCAGATGTGA (SEQ ID NO: 42)~~
- 3) AACACGTCATGTGCAGGAGTGAC;  
CGGGTGACAGTTGTGATCTAAGG (SEQ ID NO:43)

30

35

In PCR techniques, oligonucleotide primers, as listed  
 above, complementary to the two 3' borders of the DNA  
 region to be amplified are synthesized. The

SUBMITTED BY PATENT

SUB  
B4  
25

SUB  
B5

SUB  
B6  
30



polymerase chain reaction is then carried out using the two primers. See *PCR Protocols: A Guide to Methods and Applications* [74]. Following PCR amplification, the PCR-amplified regions of a viral DNA can be tested for their ability to hybridize to the three specific nucleic acid probes listed above. Alternatively, hybridization of a viral DNA to the above nucleic acid probes can be performed by a Southern blot procedure without viral DNA amplification and under stringent hybridization conditions as described herein.

Oligonucleotides for use as probes or PCR primers are chemically synthesized according to the solid phase phosphoramidite triester method first described by Beaucage and Carruthers [19] using an automated synthesizer, as described in Needham-VanDevanter [69]. Purification of oligonucleotides is by either native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson, J.D. and Regnier, F.E. [75A]. The sequence of the synthetic oligonucleotide can be verified using the chemical degradation method of Maxam, A.M. and Gilbert, W. [63].

25           B.   Isolation and propagation of KS-inducing strains of the Human Herpesvirus

Using conventional methods, the human herpesvirus can be propagated in vitro. For example, standard techniques for growing herpes viruses are described in Ablashi, D.V. [1]. Briefly, PHA stimulated cord blood mononuclear cells, macrophage, neuronal, or glial cell lines are cocultivated with cerebrospinal fluid, plasma, peripheral blood leukocytes, or tissue extracts containing viral infected cells or purified virus. The recipient cells are treated with 5  $\mu$ g/ml polybrene for 2 hours at 37° C prior to infection.

5 Infected cells are observed by demonstrating morphological changes, as well as being positive for antigens from the human herpesvirus by using monoclonal antibodies immunoreactive with the human herpes virus in an immunofluorescence assay.

10 For virus isolation, the virus is either harvested directly from the culture fluid by direct centrifugation, or the infected cells are harvested, homogenized or lysed and the virus is separated from cellular debris and purified by standard methods of isopycnic sucrose density gradient centrifugation.

15 One skilled in the art may isolate and propagate the DNA herpesvirus associated with Kaposi's sarcoma (KSHV) employing the following protocol. Long-term establishment of a B lymphoid cell line infected with the KSHV from body-cavity based lymphomas (RCC-1 or BHL-6) is prepared extracting DNA from the Lymphoma tissue using standard techniques [27, 49, 66].

20 The KS associated herpesvirus may be isolated from the cell DNA in the following manner. An infected cell line (BHL-6 RCC-1), which can be lysed using standard methods such as hypotonic shocking and Dounce homogenization, is first pelleted at 2000xg for 10 minutes, the supernatant is removed and centrifuged again at 10,000xg for 15 minutes to remove nuclei and organelles. The supernatant is filtered through a 0.45μ filter and centrifuged again at 100,000xg for 1 hour to pellet the virus. The virus can then be washed and centrifuged again at 100,000xg for 1 hour.

35 The DNA is tested for the presence of the KSHV by Southern blotting and PCR using the specific probes as described hereinafter. Fresh lymphoma tissue containing viable infected cells is simultaneously

SUB B7

SUB B8

filtered to form a single cell suspension by standard techniques [49, 66]. The cells are separated by standard Ficoll-Plaque centrifugation and lymphocyte layer is removed. The lymphocytes are then placed at  
5 >1x10<sup>6</sup> cells/ml into standard lymphocyte tissue culture medium, such as RMP 1640 supplemented with 10% fetal calf serum. Immortalized lymphocytes containing the KSHV virus are indefinitely grown in the culture media while nonimmortalized cells die during course of  
10 prolonged cultivation.

Further, the virus may be propagated in a new cell line by removing media supernatant containing the virus from a continuously infected cell line at a concentration of  
15 >1x10<sup>6</sup> cells/ml. The media is centrifuged at 2000xg for 10 minutes and filtered through a 0.45 $\mu$  filter to remove cells. The media is applied in a 1:1 volume with cells growing at >1x10<sup>6</sup> cells/ml for 48 hours. The cells are washed and pelleted and placed in fresh  
20 culture medium, and tested after 14 days of growth.

RCC-1 and RCC-1<sub>2FS</sub> were deposited on October 19, 1994 under ATCC Accession No. CRL 11734 and CRL 11735, respectively, pursuant to the Budapest Treaty on the  
25 International Deposit of Microorganisms for the Purposes of Patent Procedure with the Patent Culture Depository of the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209 U.S.A.

HBL-6 was deposited on November 18, 1994 under ATCC Accession No. CRL 11762 pursuant to the Budapest Treaty on the International Deposit of Microorganisms for the  
30 Purposes of Patent Procedure with the Patent Culture Depository of the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209 U.S.A.  
35

### C. Immunological Identity of the Virus

5 The KS-associated human herpesvirus can also be described immunologically. KS-associated human herpesviruses are selectively immunoreactive to antisera generated against a defined immunogen such as the viral major capsid protein depicted in Seq. ID No. 12, herein. Immunoreactivity is determined in an immunoassay using a polyclonal antiserum which was raised to the protein which is encoded by the amino acid sequence or nucleic acid sequence of SEQ ID NOS: 18-20. This antiserum is selected to have low crossreactivity against other herpes viruses and any such crossreactivity is removed by immunoabsorbtion prior to use in the immunoassay.

10 In order to produce antisera for use in an immunoassay, the protein which is encoded by the amino acid sequence or nucleic acid of SEQ ID NOS: 18-20 is isolated as described herein. For example, recombinant protein can be produced in a mammalian cell line. An inbred strain of mice such as balb/c is immunized with the protein which is encoded by the amino acid sequence or nucleic acid of SEQ ID NOS: 2-37 using a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see [32], *supra*). Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used an immunogen. Polyclonal sera are collected and titered against the immunogen protein in an immunoassay, for example, a solid phase immunoassay with the immunogen immobilized on a solid support. Polyclonal antisera with a titer of  $10^4$  or greater are selected and tested for their cross reactivity against other viruses of the gammaherpesvirinae subfamily, particularly human herpes virus types 1-7, by using a standard

immunoassay as described in [32], *supra*. These other gammaherpesvirinae virus can be isolated by standard techniques for isolation herpes viruses as described herein.

5

The ability of the above viruses to compete with the binding of the antisera to the immunogen protein is determined. The percent crossreactivity for other viruses is calculated, using standard calculations. Those antisera with less than 10% crossreactivity with each of the other viruses listed above is selected and pooled. The cross-reacting antibodies are then removed from the pooled antisera by immunoabsorption with the above-listed viruses.

15

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay procedure as described above to compare an unknown virus preparation to the specific KS herpesvirus preparation described herein and containing the nucleic acid sequence described in SEQ ID NOs: 2-37. In order to make this comparison, the immunogen protein which is encoded by the amino acid sequence or nucleic acid of SEQ ID NOs: 2-37 is the labeled antigen and the virus preparations are each assayed at a wide range of concentrations. The amount of each virus preparation required to inhibit 50% of the binding of the antisera to the labeled immunogen protein is determined. Those viruses that specifically bind to an antibody generated to an immunogen consisting of the protein of SEQ ID NOs: 2-37 are those virus where the amount of virus needed to inhibit 50% of the binding to the protein does not exceed an established amount. This amount is no more than 10 times the amount of the virus that is needed for 50% inhibition for the KS-associated herpesvirus containing the DNA sequence of SEQ ID NO: 1. Thus, the KS-associated herpesviruses

20

25

30

35

of the invention can be defined by immunological comparison to the specific strain of the KS-associated herpesvirus for which nucleic acid sequences are provided herein.

5

This invention provides, a nucleic acid molecule of at least 14 nucleotides capable of specifically hybridizing with the isolated DNA molecule. In one embodiment, the molecule is DNA. In another embodiment, the molecule is RNA. In another embodiment the nucleic acid molecule may be 14-20 nucleotides in length. In another embodiment the nucleic acid molecule may be 16 nucleotides in length.

10

This invention provides, a nucleic acid molecule of at least 14 nucleotides capable of specifically hybridizing with a nucleic acid molecule which is complementary to the isolated DNA molecule. In one embodiment, the molecule is DNA. In another embodiment, the molecule is RNA.

15

The nucleic acid molecule of at least 14 nucleotides may hybridize with moderate stringency to at least a portion of a nucleic acid molecule with a sequence shown in Figures 3A-3F (SEQ ID NOs: 1, 10-17, and 38-40).

20

High stringent hybridization conditions are selected at about 5° C lower than the thermal melting point (T<sub>m</sub>) for the specific sequence at a defined ionic strength and pH. The T<sub>m</sub> is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is at least about 0.02 molar at pH 7 and the temperature is at least about 60°C. As other factors may significantly affect the

25

30

35

stringency of hybridization, including, among others, base composition and size of the complementary strands, the presence of organic solvents, ie. salt or formamide concentration, and the extent of base mismatching, the combination of parameters is more important than the absolute measure of any one. For Example high stringency may be attained for example by overnight hybridization at about 68°C in a 6x SSC solution, washing at room temperature with 6x SSC solution, followed by washing at about 68°C in a 6x SSC in a 0.6x SSX solution.

Hybridization with moderate stringency may be attained for example by: 1) filter pre-hybridizing and hybridizing with a solution of 3x sodium chloride, sodium citrate (SSC), 50% formamide, 0.1M Tris buffer at Ph 7.5, 5x Denhardt's solution; 2.) pre-hybridization at 37°C for 4 hours; 3) hybridization at 37°C with amount of labelled probe equal to 3,000,000 cpm total for 16 hours; 4) wash in 2x SSC and 0.1% SDS solution; 5) wash 4x for 1 minute each at room temperature at 4x at 60°C for 30 minutes each; and 6) dry and expose to film.

The phrase "selectively hybridizing to" refers to a nucleic acid probe that hybridizes, duplexes or binds only to a particular target DNA or RNA sequence when the target sequences are present in a preparation of total cellular DNA or RNA. By selectively hybridizing it is meant that a probe binds to a given target in a manner that is detectable in a different manner from non-target sequence under high stringency conditions of hybridization. in a different "Complementary" or "target" nucleic acid sequences refer to those nucleic acid sequences which selectively hybridize to a nucleic acid probe. Proper annealing conditions depend, for example, upon a probe's length, base

composition, and the number of mismatches and their position on the probe, and must often be determined empirically. For discussions of nucleic acid probe design and annealing conditions, see, for example, 5 Sambrook et al., [81] or Ausubel, F., et al., [8].

It will be readily understood by those skilled in the art and it is intended here, that when reference is made to particular sequence listings, such reference 10 includes sequences which substantially correspond to its complementary sequence and those described including allowances for minor sequencing errors, single base changes, deletions, substitutions and the like, such that any such sequence variation 15 corresponds to the nucleic acid sequence of the pathogenic organism or disease marker to which the relevant sequence listing relates.

Nucleic acid probe technology is well known to those 20 skilled in the art who readily appreciate that such probes may vary greatly in length and may be labeled with a detectable label, such as a radioisotope or fluorescent dye, to facilitate detection of the probe. DNA probe molecules may be produced by insertion of a 25 DNA molecule having the full-length or a fragment of the isolated nucleic acid molecule of the DNA virus into suitable vectors, such as plasmids or bacteriophages, followed by transforming into suitable bacterial host cells, replication in the transformed 30 bacterial host cells and harvesting of the DNA probes, using methods well known in the art. Alternatively, probes may be generated chemically from DNA synthesizers.

35 DNA virus nucleic acid rearrangements/mutations may be detected by Southern blotting, single stranded conformational polymorphism gel electrophoresis



(SSCP), PCR or other DNA based techniques, or for RNA species by Northern blotting, PCR or other RNA-based techniques.

- 5 RNA probes may be generated by inserting the full  
length or a fragment of the isolated nucleic acid  
molecule of the DNA virus downstream of a  
bacteriophage promoter such as T3, T7 or SP6. Large  
amounts of RNA probe may be produced by incubating the  
10 labeled nucleotides with a linearized isolated nucleic  
acid molecule of the DNA virus or its fragment where  
it contains an upstream promoter in the presence of  
the appropriate RNA polymerase.
- 15 As defined herein nucleic acid probes may be DNA or  
RNA fragments. DNA fragments can be prepared, for  
example, by digesting plasmid DNA, or by use of PCR,  
or synthesized by either the phosphoramidite method  
described by Beaucage and Carruthers, [19], or by the  
20 triester method according to Matteucci, et al., [62],  
both incorporated herein by reference. A double  
stranded fragment may then be obtained, if desired, by  
annealing the chemically synthesized single strands  
together under appropriate conditions or by  
25 synthesizing the complementary strand using DNA  
polymerase with an appropriate primer sequence. Where  
a specific sequence for a nucleic acid probe is given,  
it is understood that the complementary strand is also  
identified and included. The complementary strand  
30 will work equally well in situations where the target  
is a double-stranded nucleic acid. It is also  
understood that when a specific sequence is identified  
for use a nucleic probe, a subsequence of the listed  
sequence which is 25 basepairs or more in length is  
35 also encompassed for use as a probe.

The DNA molecules of the subject invention also include DNA molecules coding for polypeptide analogs, fragments or derivatives of antigenic polypeptides which differ from naturally-occurring forms in terms of the identity or location of one or more amino acid residues (deletion analogs containing less than all of the residues specified for the protein, substitution analogs wherein one or more residues specified are replaced by other residues and addition analogs where in one or more amino acid residues is added to a terminal or medial portion of the polypeptides) and which share some or all properties of naturally-occurring forms. These molecules include: the incorporation of codons "preferred" for expression by selected non-mammalian hosts; the provision of sites for cleavage by restriction endonuclease enzymes; and the provision of additional initial, terminal or intermediate DNA sequences that facilitate construction of readily expressed vectors.

This invention provides for an isolated DNA molecule which encodes at least a portion of a Kaposi's sarcoma associated herpesvirus: virion polypeptide 23, major capsid protein, capsid proteins, thymidine kinase, or tegument protein.

This invention also provides a method of producing a polypeptide encoded by isolated DNA molecule, which comprises growing the above host vector system under suitable conditions permitting production of the polypeptide and recovering the polypeptide so produced.

This invention provides an isolated peptide encoded by the isolated DNA molecule associated with Kaposi's sarcoma. In one embodiment the peptide may be a polypeptide. Further, this invention provides a host

cell which expresses the polypeptide of isolated DNA molecule.

5 In one embodiment the isolated peptide or polypeptide is encoded by at least a portion of an isolated DNA molecule. In another embodiment the isolated peptide or polypeptide is encoded by at least a portion of a nucleic acid molecule with a sequence as set forth in (SEQ ID NOs: 2-37).

10 Further, the isolated peptide or polypeptide encoded by the isolated DNA molecule may be linked to a second nucleic acid molecule to form a fusion protein by expression in a suitable host cell. In one embodiment the second nucleic acid molecule encodes beta-galactosidase. Other nucleic acid molecules which are used to form a fusion protein are known to those skilled in the art.

20 This invention provides an antibody which specifically binds to the peptide or polypeptide encoded by the isolated DNA molecule. In one embodiment the antibody is a monoclonal antibody. In another embodiment the antibody is a polyclonal antibody.

25 The antibody or DNA molecule may be labelled with a detectable marker including, but not limited to: a radioactive label, or a colorimetric, a luminescent, or a fluorescent marker, or gold. Radioactive labels include, but are not limited to:  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{33}\text{P}$ ;  $^{35}\text{S}$ ,  $^{36}\text{Cl}$ ,  $^{51}\text{Cr}$ ,  $^{57}\text{Co}$ ,  $^{59}\text{Co}$ ,  $^{59}\text{Fe}$ ,  $^{90}\text{Y}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ , and  $^{186}\text{Re}$ . Fluorescent markers include but are not limited to: fluorescein, rhodamine and auramine. Colorimetric markers include, but are not limited to: biotin, and digoxigenin. Methods of producing the polyclonal or  
30 monoclonal antibody are known to those of ordinary skill in the art.  
35

Further, the antibody or nucleic acid molecule complex may be detected by a second antibody which may be linked to an enzyme, such as alkaline phosphatase or horseradish peroxidase. Other enzymes which may be employed are well known to one of ordinary skill in the art.

This invention provides a method to select specific regions on the polypeptide encoded by the isolated DNA molecule of the DNA virus to generate antibodies. The protein sequence may be determined from the cDNA sequence. Amino acid sequences may be analyzed by methods well known to those skilled in the art to determine whether they produce hydrophobic or hydrophilic regions in the proteins which they build. In the case of cell membrane proteins, hydrophobic regions are well known to form the part of the protein that is inserted into the lipid bilayer of the cell membrane, while hydrophilic regions are located on the cell surface, in an aqueous environment. Usually, the hydrophilic regions will be more immunogenic than the hydrophobic regions. Therefore the hydrophilic amino acid sequences may be selected and used to generate antibodies specific to polypeptide encoded by the isolated nucleic acid molecule encoding the DNA virus. The selected peptides may be prepared using commercially available machines. As an alternative, DNA, such as a cDNA or a fragment thereof, may be cloned and expressed and the resulting polypeptide recovered and used as an immunogen.

Polyclonal antibodies against these peptides may be produced by immunizing animals using the selected peptides. Monoclonal antibodies are prepared using hybridoma technology by fusing antibody producing B cells from immunized animals with myeloma cells and selecting the resulting hybridoma cell line producing

the desired antibody. Alternatively, monoclonal antibodies may be produced by in vitro techniques known to a person of ordinary skill in the art. These antibodies are useful to detect the expression of polypeptide encoded by the isolated DNA molecule of the DNA virus in living animals, in humans, or in biological tissues or fluids isolated from animals or humans.

10 II. Immunoassays

The antibodies raised against the viral strain or peptides may be detectably labelled, utilizing conventional labelling techniques well-known to the art. Thus, the antibodies may be radiolabelled using, for example, radioactive isotopes such as  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ , and  $^{35}\text{S}$ .

The antibodies may also be labelled using fluorescent labels, enzyme labels, free radical labels, or bacteriophage labels, using techniques known in the art. Typical fluorescent labels include fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, and Texas Red.

Since specific enzymes may be coupled to other molecules by covalent links, the possibility also exists that they might be used as labels for the production of tracer materials. Suitable enzymes include alkaline phosphatase, beta-galactosidase, glucose-6-phosphate dehydrogenase, maleate dehydrogenase, and peroxidase. Two principal types of enzyme immunoassay are the enzyme-linked immunosorbent assay (ELISA), and the homogeneous enzyme immunoassay, also known as enzyme-multiplied immunoassay (EMIT, Syva Corporation, Palo Alto, CA). In the ELISA system, separation may be achieved, for example, by

the use of antibodies coupled to a solid phase. The EMIT system depends on deactivation of the enzyme in the tracer-antibody complex; the activity can thus be measured without the need for a separation step.

5

Additionally, chemiluminescent compounds may be used as labels. Typical chemiluminescent compounds include luminol, isoluminol, aromatic acridinium esters, imidazoles, acridinium salts, and oxalate esters. Similarly, bioluminescent compounds may be utilized for labelling, the bioluminescent compounds including luciferin, luciferase, and aequorin.

10

Once labeled, the antibody may be employed to identify and quantify immunologic counterparts (antibody or antigenic polypeptide) utilizing techniques well-known to the art.

15

A description of a radioimmunoassay (RIA) may be found in *Laboratory Techniques in Biochemistry and Molecular Biology* [52], with particular reference to the chapter entitled "An Introduction to Radioimmune Assay and Related Techniques" by Chard, T., incorporated by reference herein.

20

25

A description of general immunometric assays of various types can be found in the following U.S. Pat. Nos. 4,376,110 (David et al.) or 4,098,876 (Piasio).

30

A. Assays for viral antigens

In addition to the detection of the causal agent using nucleic acid hybridization technology, one can use immunoassays to detect for the virus, specific peptides, or for antibodies to the virus or peptides. A general overview of the applicable technology is in

35

Harlow and Lane [32], incorporated by reference herein.

5 In one embodiment, antibodies to the human herpesvirus can be used to detect the agent in the sample. In brief, to produce antibodies to the agent or peptides, the sequence being targeted is expressed in transfected cells, preferably bacterial cells, and purified. The product is injected into a mammal capable of producing antibodies. Either monoclonal or polyclonal antibodies (as well as any recombinant antibodies) specific for the gene product can be used in various immunoassays. Such assays include competitive immunoassays, radioimmunoassays, Western blots, ELISA, indirect immunofluorescent assays and the like. For competitive immunoassays, see Harlow and Lane [32] at pages 567-573 and 584-589.

10  
15  
20 Monoclonal antibodies or recombinant antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells or other lymphocytes from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell (see, Kohler and Milstein [50], incorporated herein by reference). Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host. New techniques using recombinant phage antibody expression systems can also be used to generate monoclonal antibodies. See for example: McCafferty, J et al.

25  
30  
35

[64]; Hoogenboom, H.R. et al. [39]; and Marks, J.D. et al. [60].

Such peptides may be produced by expressing the  
5 specific sequence in a recombinantly engineered cell  
such as bacteria, yeast, filamentous fungal, insect  
(especially employing baculoviral vectors), and  
mammalian cells. Those of skill in the art are  
knowledgeable in the numerous expression systems  
10 available for expression of herpes virus protein.

Briefly, the expression of natural or synthetic  
nucleic acids encoding viral protein will typically be  
achieved by operably linking the desired sequence or  
15 portion thereof to a promoter (which is either  
constitutive or inducible), and incorporated into an  
expression vector. The vectors are suitable for  
replication or integration in either prokaryotes or  
eukaryotes. Typical cloning vectors contain  
20 antibiotic resistance markers, genes for selection of  
transformants, inducible or regulatable promoter  
regions, and translation terminators that are useful  
for the expression of viral genes.

25 Methods for the expression of cloned genes in bacteria  
are also well known. In general, to obtain high level  
expression of a cloned gene in a prokaryotic system,  
it is advisable to construct expression vectors  
containing a strong promoter to direct mRNA  
30 transcription. The inclusion of selection markers in  
DNA vectors transformed in *E. coli* is also useful.  
Examples of such markers include genes specifying  
resistance to antibiotics. See [81] *supra*, for  
details concerning selection markers and promoters for  
35 use in *E. coli*. Suitable eukaryote hosts may include  
plant cells, insect cells, mammalian cells, yeast, and  
filamentous fungi.



Methods for characterizing naturally processed peptides bound to MHC (major histocompatibility complex) I molecules have been developed. See, Falk et al. [24], and PCT publication No. WO 92/21033 published November 26, 1992, both of which are incorporated by reference herein. Typically, these methods involve isolation of MHC class I molecules by immunoprecipitation or affinity chromatography from an appropriate cell or cell line. Other methods involve direct amino acid sequencing of the more abundant peptides in various HPLC fractions by known automatic sequencing of peptides eluted from Class I molecules of the B cell type (Jardetzkey, et al. [45], incorporated by reference herein, and of the human MHC class I molecule, HLA-A2.1 type by mass spectrometry (Hunt, et al. [40], incorporated by reference herein). See also, Röttschke and Falk [79], incorporated by reference herein for a general review of the characterization of naturally processed peptides in MHC class I. Further, Marloes, et al. [61], incorporated by reference herein, describe how class I binding motifs can be applied to the identification of potential viral immunogenic peptides in vitro.

The peptides described herein produced by recombinant technology may be purified by standard techniques well known to those of skill in the art. Recombinantly produced viral sequences can be directly expressed or expressed as a fusion protein. The protein is then purified by a combination of cell lysis (e.g., sonication) and affinity chromatography. For fusion products, subsequent digestion of the fusion protein with an appropriate proteolytic enzyme releases the desired peptide.

The proteins may be purified to substantial purity by standard techniques well known in the art, including

selective precipitation with such substances as ammonium sulfate, column chromatography, immunopurification methods, and others. See, for instance, Scopes, R. [84], incorporated herein by  
5 reference.

B. Serological tests for the presence of antibodies to the human herpesvirus.

10 This invention further embraces diagnostic kits for detecting the presence of a KS agent in biological samples, such as serum or solid tissue samples, comprising a container containing antibodies to the human herpesvirus, and instructional material for  
15 performing the test. Alternatively, inactivated viral particles or peptides or viral proteins derived from the human herpesvirus may be used in a diagnostic kit to detect for antibodies specific to the KS associated human herpesvirus.

20 Diagnostic kits for detecting the presence of a KS agent in tissue samples, such as skin samples or samples of other affected tissue, comprising a container containing a nucleic acid sequence specific for the human herpesvirus and instructional material  
25 for detecting the KS-associated herpesvirus are also included. A container containing nucleic acid primers to any one of such sequences is optionally included as are antibodies to the human herpesvirus as described  
30 herein.

Antibodies reactive with antigens of the human herpesvirus can also be measured by a variety of immunoassay methods that are similar to the procedures  
35 described above for measurement of antigens. For a review of immunological and immunoassay procedures applicable to the measurement of antibodies by

immunoassay techniques, see *Basic and Clinical Immunology* 7th Edition [12], and [32], *supra*.

5 In brief, immunoassays to measure antibodies reactive  
with antigens of the KS-associated human herpesvirus  
can be either competitive or noncompetitive binding  
assays. In competitive binding assays, the sample  
analyte competes with a labeled analyte for specific  
10 binding sites on a capture agent bound to a solid  
surface. Preferably the capture agent is a purified  
recombinant human herpesvirus protein produced as  
described above. Other sources of human herpesvirus  
proteins, including isolated or partially purified  
15 naturally occurring protein, may also be used.  
Noncompetitive assays are typically sandwich assays,  
in which the sample analyte is bound between two  
analyte-specific binding reagents. One of the binding  
agents is used as a capture agent and is bound to a  
solid surface. The second binding agent is labelled  
20 and is used to measure or detect the resultant complex  
by visual or instrument means. A number of  
combinations of capture agent and labelled binding  
agent can be used. A variety of different immunoassay  
formats, separation techniques and labels can be also  
25 be used similar to those described above for the  
measurement of the human herpesvirus antigens.

Hemagglutination Inhibition (HI) and Complement  
30 Fixation (CF) which are two laboratory tests that can  
be used to detect infection with human herpesvirus by  
testing for the presence of antibodies against the  
virus or antigens of the virus.

Serological methods can be also be useful when one  
35 wishes to detect antibody to a specific variant. For  
example, one may wish to see how well a vaccine  
recipient has responded to the new variant.

Alternatively, one may take serum from a patient to see which variant the patient responds to the best.

5 This invention provides an antagonist capable of blocking the expression of the peptide or polypeptide encoded by the isolated DNA molecule. In one embodiment the antagonist is capable of hybridizing with a double stranded DNA molecule. In another embodiment the antagonist is a triplex oligonucleotide  
10 capable of hybridizing to the DNA molecule. In another embodiment the triplex oligonucleotide is capable of binding to at least a portion of the isolated DNA molecule with a nucleotide sequence as shown in Figure 3A-3F (SEQ ID NOs: 1-37).

15 This invention provides an antisense molecule capable of hybridizing to the isolated DNA molecule. In one embodiment the antisense molecule is DNA. In another embodiment the antisense molecule is RNA.

20 The antisense molecule may be DNA or RNA or variants thereof (i.e. DNA or RNA with a protein backbone). The present invention extends to the preparation of antisense nucleotides and ribozymes that may be used  
25 to interfere with the expression of the receptor recognition proteins at the translation of a specific mRNA, either by masking that MRNA with an antisense nucleic acid or cleaving it with a ribozyme.

30 Antisense nucleic acids are DNA or RNA molecules that are complementary to at least a portion of a specific MRNA molecule. In the cell, they hybridize to that MRNA, forming a double stranded molecule. The cell  
35 does not translate an MRNA in this double-stranded form. Therefore, antisense nucleic acids interfere with the expression of MRNA into protein. Oligomers of about fifteen nucleotides and molecules that

hybridize to the AUG initiation codon are particularly efficient, since they are easy to synthesize and are likely to pose fewer problems than larger molecules upon introduction to cells.

5

This invention provides a transgenic nonhuman mammal which comprises at least a portion of the isolated DNA molecule introduced into the mammal at an embryonic stage. Methods of producing a transgenic nonhuman mammal are known to those skilled in the art.

10

This invention provides a cell line containing the isolated KS associated herpesvirus of the subject invention. In one embodiment the isolated DNA molecule is artificially introduced into the cell. Cell lines include, but are not limited to: fibroblasts, such as HFF, NIH/3T3; Epithelial cells, such as 5637; lymphocytes, such as FCB; T-cells, such as CCRF-CEM (ATCC CCL 119); B-cells, such as BJAB and Raji (ATCC CCL 86); and myeloid cells such as K562 (ATCC CCL 243); Vero cells and carcinoma cells. Methods of producing such cell lines are known to those skilled in the art. In one embodiment the isolated KS associated herpesvirus is introduced into a RCC-1 cell line.

15

20

25

III. In vitro diagnostic assays for the detection of  
KS

30

35

This invention provides a method of diagnosing Kaposi's sarcoma in a subject which comprises: (a) obtaining a nucleic acid molecule from a tumor lesion of the subject; (b) contacting the nucleic acid molecule with a labelled nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with the isolated DNA, under hybridizing conditions; and (c) determining the presence of the

nucleic acid molecule hybridized, the presence of which is indicative of Kaposi's sarcoma in the subject, thereby diagnosing Kaposi's sarcoma in the subject.

5

In one embodiment the DNA molecule from the tumor lesion is amplified before step (b). In another embodiment PCR is employed to amplify the nucleic acid molecule. Methods of amplifying nucleic acid molecules are known to those skilled in the art.

10

A person of ordinary skill in the art will be able to obtain appropriate DNA sample for diagnosing Kaposi's sarcoma in the subject. The DNA sample obtained by the above described method may be cleaved by restriction enzyme. The uses of restriction enzymes to cleave DNA and the conditions to perform such cleavage are well-known in the art.

15

20

In the above described methods, a size fractionation may be employed which is effected by a polyacrylamide gel. In one embodiment, the size fractionation is effected by an agarose gel. Further, transferring the DNA fragments into a solid matrix may be employed before a hybridization step. One example of such solid matrix is nitrocellulose paper.

25

This invention provides a method of diagnosing Kaposi's sarcoma in a subject which comprises: (a) obtaining a nucleic acid molecule from a suitable bodily fluid of the subject; (b) contacting the nucleic acid molecule with a labelled nucleic acid molecules of at least 15 nucleotides capable of specifically hybridizing with the isolated DNA, under hybridizing conditions; and (c) determining the presence of the nucleic acid molecule hybridized, the

35

presence of which is indicative of Kaposi's sarcoma in the subject, thereby diagnosing Kaposi's sarcoma in the subject.

5 This invention provides a method of diagnosing a DNA virus in a subject, which comprises (a) obtaining a suitable bodily fluid sample from the subject, (b) contacting the suitable bodily fluid of the subject to a support having already bound thereto a Kaposi's sarcoma antibody, so as to bind the Kaposi's sarcoma antibody to a specific Kaposi's sarcoma antigen, (c) removing unbound bodily fluid from the support, and (d) determining the level of Kaposi's sarcoma antibody bound by the Kaposi's sarcoma antigen, thereby diagnosing the subject for Kaposi's sarcoma.

20 This invention provides a method of diagnosing Kaposi's sarcoma in a subject, which comprises (a) obtaining a suitable bodily fluid sample from the subject, (b) contacting the suitable bodily fluid of the subject to a support having already bound thereto a Kaposi's sarcoma antigen, so as to bind Kaposi's sarcoma antigen to a specific Kaposi's sarcoma antibody, (c) removing unbound bodily fluid from the support, and (d) determining the level of the Kaposi's sarcoma antigen bound by the Kaposi's sarcoma antibody, thereby diagnosing Kaposi's sarcoma.

30 This invention provides a method of detecting expression of a DNA virus associated with Kaposi's sarcoma in a cell which comprises obtaining total cDNA obtained from the cell, contacting the cDNA so obtained with a labelled DNA molecule under hybridizing conditions, determining the presence of cDNA hybridized to the molecule, and thereby detecting the expression of the DNA virus. In one embodiment

mRNA is obtained from the cell to detect expression of the DNA virus.

5 The suitable bodily fluid sample is any bodily fluid sample which would contain Kaposi's sarcoma antibody, antigen or fragments thereof. A suitable bodily fluid includes, but is not limited to: serum, plasma, cerebrospinal fluid, lymphocytes, urine, transudates, or exudates. In the preferred embodiment, the  
10 suitable bodily fluid sample is serum or plasma. In addition, the bodily fluid sample may be cells from bone marrow, or a supernatant from a cell culture. Methods of obtaining a suitable bodily fluid sample from a subject are known to those skilled in the art. Methods of determining the level of antibody or  
15 antigen include, but are not limited to: ELISA, IFA, and Western blotting. Other methods are known to those skilled in the art. Further, a subject infected with a DNA virus associated with Kaposi's sarcoma may  
20 be diagnosed with the above described methods.

The detection of the human herpesvirus and the detection of virus-associated KS are essentially identical processes. The basic principle is to detect  
25 the virus using specific ligands that bind to the virus but not to other proteins or nucleic acids in a normal human cell or its environs. The ligands can either be nucleic acid or antibodies. The ligands can be naturally occurring or genetically or physically  
30 modified such as nucleic acids with non-natural or antibody derivatives, i.e., Fab or chimeric antibodies. Serological tests for detection of antibodies to the virus may also be performed by using protein antigens obtained from the human herpesvirus,  
35 and described herein.



Samples can be taken from patients with KS or from patients at risk for KS, such as AIDS patients. Typically the samples are taken from blood (cells, serum and/or plasma) or from solid tissue samples such as skin lesions. The most accurate diagnosis for KS will occur if elevated titers of the virus are detected in the blood or in involved lesions. KS may also be indicated if antibodies to the virus are detected and if other diagnostic factors for KS is present.

A. Nucleic acid assays.

The diagnostic assays of the invention can be nucleic acid assays such as nucleic acid hybridization assays and assays which detect amplification of specific nucleic acid to detect for a nucleic acid sequence of the human herpesvirus described herein.

Accepted means for conducting hybridization assays are known and general overviews of the technology can be had from a review of: *Nucleic Acid Hybridization: A Practical Approach* [72]; *Hybridization of Nucleic Acids Immobilized on Solid Supports* [41]; *Analytical Biochemistry* [4] and Innis et al., *PCR Protocols* [74], supra, all of which are incorporated by reference herein.

If PCR is used in conjunction with nucleic acid hybridization, primers are designed to target a specific portion of the nucleic acid of the herpesvirus. For example, the primers set forth in SEQ ID NOs: 38-40 may be used to target detection of regions of the herpesvirus genome encoding ORF 25 homologue - ORF 32 homologue. From the information provided herein, those of skill in the art will be able to select appropriate specific primers.

Target specific probes may be used in the nucleic acid hybridization diagnostic assays for KS. The probes are specific for or complementary to the target of interest. For precise allelic differentiations, the probes should be about 14 nucleotides long and preferably about 20-30 nucleotides. For more general detection of the human herpesvirus of the invention, nucleic acid probes are about 50 to about 1000 nucleotides, most preferably about 200 to about 400 nucleotides.

A sequence is "specific" for a target organism of interest if it includes a nucleic acid sequence which when detected is determinative of the presence of the organism in the presence of a heterogeneous population of proteins and other biologics. A specific nucleic acid probe is targeted to that portion of the sequence which is determinative of the organism and will not hybridize to other sequences especially those of the host where a pathogen is being detected.

The specific nucleic acid probe can be RNA or DNA polynucleotide or oligonucleotide, or their analogs. The probes may be single or double stranded nucleotides. The probes of the invention may be synthesized enzymatically, using methods well known in the art (e.g., nick translation, primer extension, reverse transcription, the polymerase chain reaction, and others) or chemically (e.g., by methods such as the phosphoramidite method described by Beaucage and Carruthers [19], or by the triester method according to Matteucci, et al. [62], both incorporated herein by reference).

The probe must be of sufficient length to be able to form a stable duplex with its target nucleic acid in the sample, i.e., at least about 14 nucleotides, and

may be longer (e.g., at least about 50 or 100 bases in length). Often the probe will be more than about 100 bases in length. For example, when probe is prepared by nick-translation of DNA in the presence of labeled nucleotides the average probe length may be about 100-  
5 600 bases.

As noted above, the probe will be capable of specific hybridization to a specific KS-associated herpes virus nucleic acid. Such "specific hybridization" occurs when a probe hybridizes to a target nucleic acid, as evidenced by a detectable signal, under conditions in which the probe does not hybridize to other nucleic acids (e.g., animal cell or other bacterial nucleic acids) present in the sample. A variety of factors including the length and base composition of the probe, the extent of base mismatching between the probe and the target nucleic acid, the presence of salt and organic solvents, probe concentration, and the temperature affect hybridization, and optimal hybridization conditions must often be determined empirically. For discussions of nucleic acid probe design and annealing conditions, see, for example, [81], supra, Ausubel, F., et al. [8] [hereinafter referred to as Sambrook], *Methods in Enzymology* [67] or *Hybridization with Nucleic Acid Probes* [42] all of which are incorporated herein by reference.  
10  
15  
20  
25

Usually, at least a part of the probe will have considerable sequence identity with the target nucleic acid. Although the extent of the sequence identity required for specific hybridization will depend on the length of the probe and the hybridization conditions, the probe will usually have at least 70% identity to the target nucleic acid, more usually at least 80%  
30 35 identity, still more usually at least 90% identity and most usually at least 95% or 100% identity.

A probe can be identified as capable of hybridizing specifically to its target nucleic acid by hybridizing the probe to a sample treated according the protocol of this invention where the sample contains both target virus and animal cells (e.g., nerve cells). A probe is specific if the probe's characteristic signal is associated with the herpesvirus DNA in the sample and not generally with the DNA of the host cells and non-biological materials (e.g., substrate) in a sample.

The following stringent hybridization and washing conditions will be adequate to distinguish a specific probe (e.g., a fluorescently labeled DNA probe) from a probe that is not specific: incubation of the probe with the sample for 12 hours at 37°C in a solution containing denatured probe, 50% formamide, 2X SSC, and 0.1% (w/v) dextran sulfate, followed by washing in 1X SSC at 70°C for 5 minutes; 2X SSC at 37°C for 5 minutes; 0.2X SSC at room temperature for 5 minutes, and H<sub>2</sub>O at room temperature for 5 minutes. Those of skill will be aware that it will often be advantageous in nucleic acid hybridizations (*i.e.*, *in situ*, Southern, or other) to include detergents (e.g., sodium dodecyl sulfate), chelating agents (e.g., EDTA) or other reagents (e.g., buffers, Denhardt's solution, dextran sulfate) in the hybridization or wash solutions. To test the specificity of the virus specific probes, the probes can be tested on host cells containing the KS-associated herpesvirus and compared with the results from cells containing non-KS-associated virus.

It will be apparent to those of ordinary skill in the art that a convenient method for determining whether a probe is specific for a KS-associated viral nucleic acid utilizes a Southern blot (or Dot blot) using DNA

prepared from one or more KS-associated human herpesviruses of the invention. Briefly, to identify a target specific probe DNA is isolated from the virus. Test DNA either viral or cellular is transferred to a solid (e.g., charged nylon) matrix. The probes are labelled following conventional methods. Following denaturation and/or prehybridization steps known in the art, the probe is hybridized to the immobilized DNAs under stringent conditions. Stringent hybridization conditions will depend on the probe used and can be estimated from the calculated  $T_m$  (melting temperature) of the hybridized probe (see, e.g., Sambrook for a description of calculation of the  $T_m$ ). For radioactively-labeled DNA or RNA probes an example of stringent hybridization conditions is hybridization in a solution containing denatured probe and 5x SSC at 65°C for 8-24 hours followed by washes in 0.1x SSC, 0.1% SDS (sodium dodecyl sulfate) at 50-65°C. In general, the temperature and salt concentration are chosen so that the post hybridization wash occurs at a temperature that is about 5°C below the  $T_m$  of the hybrid. Thus for a particular salt concentration the temperature may be selected that is 5°C below the  $T_m$  or conversely, for a particular temperature, the salt concentration is chosen to provide a  $T_m$  for the hybrid that is 5°C warmer than the wash temperature. Following stringent hybridization and washing, a probe that hybridizes to the KS-associated viral DNA but not to the non-KS associated viral DNA, as evidenced by the presence of a signal associated with the appropriate target and the absence of a signal from the non-target nucleic acids, is identified as specific for the KS associated virus. It is further appreciated that in determining probe specificity and in utilizing the method of this invention to detect KS-associated herpesvirus, a certain amount of background signal is typical and can

easily be distinguished by one of skill from a specific signal. Two fold signal over background is acceptable.

5 A preferred method for detecting the KS-associated herpesvirus is the use of PCR and/or dot blot hybridization. The presence or absence of an KS agent for detection or prognosis, or risk assessment for KS includes Southern transfers, solution hybridization or  
10 non-radioactive detection systems, all of which are well known to those of skill in the art. Hybridization is carried out using probes. Visualization of the hybridized portions allows the qualitative determination of the presence or absence  
15 of the causal agent.

Similarly, a Northern transfer may be used for the detection of message in samples of RNA or reverse transcriptase PCR and cDNA can be detected by methods  
20 described above. This procedure is also well known in the art. See [81] incorporated by reference herein.

An alternative means for determining the presence of the human herpesvirus is in situ hybridization, or  
25 more recently, in situ polymerase chain reaction. In situ PCR is described in Neuvo et al. [71], Intracellular localization of polymerase chain reaction (PCR)-amplified Hepatitis C cDNA; Bagasra et al. [10], Detection of Human Immunodeficiency virus  
30 type 1 provirus in mononuclear cells by in situ polymerase chain reaction; and Heniford et al. [35], Variation in cellular EGF receptor mRNA expression demonstrated by in situ reverse transcriptase polymerase chain reaction. In situ hybridization  
35 assays are well known and are generally described in *Methods Enzymol.* [67] incorporated by reference herein. In an in situ hybridization, cells are fixed

to a solid support, typically a glass slide. The cells are then contacted with a hybridization solution at a moderate temperature to permit annealing of target-specific probes that are labelled. The probes  
5 are preferably labelled with radioisotopes or fluorescent reporters.

The above described probes are also useful for in-situ hybridization or in order to locate tissues which express this gene, or for other hybridization assays for the presence of this gene or its mRNA in various biological tissues. In-situ hybridization is a sensitive localization method which is not dependent on expression of antigens or native vs. denatured  
10 conditions.

Oligonucleotide (oligo) probes, synthetic oligonucleotide probes or riboprobes made from KSHV phagemids/plasmids, are relatively homogeneous reagents and successful hybridization conditions in tissue sections is readily transferable from one probe to another. Commercially synthesized oligonucleotide probes are prepared against the identified genes. These probes are chosen for length (45-65 mers), high  
20 G-C content (50-70%) and are screened for uniqueness against other viral sequences in GenBank.

Oligonucleotides are 3' end-labeled with [ $\alpha$ -<sup>35</sup>S]dATP to specific activities in the range of  $1 \times 10^{10}$  dpm/ug using terminal deoxynucleotidyl transferase. Unincorporated labeled nucleotides are removed from the oligo probe by centrifugation through a Sephadex G-25 column or by elution from a Waters Sep Pak C-18 column.  
30

35

KS tissue embedded in OCT compound and snap frozen in freezing isopentane cooled with dry ice is cut at 6  $\mu$ m

intervals and thawed onto 3-aminopropyltriethoxysilane treated slides and allowed to air dry. The slides are then be fixed in 4% freshly prepared paraformaldehyde, rinsed in water. Formalin-fixed, paraffin embedded KS tissues cut at 6  $\mu$ m and baked onto glass slides can also be used. The sections are then deparaffinized in xylenes and rehydrated through graded alcohols. Prehybridization in 20mM Tris Ph 7.5, 0.02% Denhardt's solution, 10% dextran sulfate for 30 min at 37°C is followed by hybridization overnight in a solution of 50% formamide (v/v), 10% dextran sulfate (w/v), 20mM sodium phosphate (Ph 7.4), 3X SSC, 1X Denhardt's solution, 100 ug/ml salmon sperm DNA, 125 ug/ml yeast tRNA and the oligo probe ( $10^6$ cpm/ml) at 42°C overnight. The slides are washed twice with 2X SSC and twice with 1X SSC for 15 minutes each at room temperature and visualized by autoradiography. Briefly, sections are dehydrated through graded alcohols containing 0.3M ammonium acetate and air dried. The slides are dipped in Kodak NTB2 emulsion, exposed for days to weeks, developed, and counterstained with hematoxylin and eoxin. Alternative immunohistochemical protocols may be employed which are known to those skilled in the art.

25

#### IV. Treatment of human herpesvirus-induced KS

This invention provides a method of treating a subject with Kaposi's sarcoma, comprising administering to the subject an effective amount of the antisense molecule capable of hybridizing to the isolated DNA molecule under conditions such that the antisense molecule selectively enters a tumor cell of the subject, so as to treat the subject.



5 This invention provides a method for treating a subject with Kaposi's sarcoma (KS) comprising administering to the subject having a human herpesvirus-associated KS a pharmaceutically effective amount of an antiviral agent in a pharmaceutically acceptable carrier, wherein the agent is effective to treat the subject with KS-associated human herpesvirus.

10 Further, this invention provides a method of prophylaxis or treatment for Kaposi's sarcoma (KS) by administering to a patient at risk for KS, an antibody that binds to the human herpesvirus in a pharmaceutically acceptable carrier. In one  
15 embodiment the antiviral drug is used to treat a subject with the DNA herpesvirus of the subject invention.

20 The use of combinations of antiviral drugs and sequential treatments are useful for treatment of herpesvirus infections and will also be useful for the treatment of herpesvirus-induced KS. For example, Snoeck et al. [88], found additive or synergistic  
25 effects against CMV when combining antiherpes drugs (e.g., combinations of zidovudine [3'-azido-3'-deoxythymidine, AZT] with HPMPC, ganciclovir, foscarnet or acyclovir or of HPMPC with other antivirals). Similarly, in treatment of  
30 cytomegalovirus retinitis, induction with ganciclovir followed by maintenance with foscarnet has been suggested as a way to maximize efficacy while minimizing the adverse side effects of either treatment alone. An anti-herpetic composition that  
35 contains acyclovir and, e.g., 2-acetylpyridine-5-((2-pyridylamino)thiocarbonyl)-thiocarbonohydrazone is described in U.S. Pat. 5,175,165 (assigned to

Burroughs Wellcome Co.). Combinations of TS-inhibitors and viral TK-inhibitors in antiherpetic medicines are disclosed in U.S. Pat. 5,137,724, assigned to Stichting Rega VZW. A synergistic  
5 inhibitory effect on EBV replication using certain ratios of combinations of HPMPC with AZT was reported by Lin et al. [56].

U.S. Patent Nos. 5,164,395 and 5,021,437 (Blumenkopf;  
10 Burroughs Wellcome) describe the use of a ribonucleotide reductase inhibitor (an acetylpyridine derivative) for treatment of herpes infections, including the use of the acetylpyridine derivative in combination with acyclovir. U.S. Patent No. 5,137,724  
15 (Balzari et al. [11]) describes the use of thymidylate synthase inhibitors (e.g., 5-fluoro-uracil and 5-fluoro-2'-deoxyuridine) in combination with compounds having viral thymidine kinase inhibiting activity.

20 With the discovery of a disease causal agent for KS now identified, effective therapeutic or prophylactic protocols to alleviate or prevent the symptoms of herpes virus-associated KS can be formulated. Due to the viral nature of the disease, antiviral agents have  
25 application here for treatment, such as interferons, nucleoside analogues, ribavirin, amantadine, and pyrophosphate analogues of phosphonoacetic acid (foscarnet) (reviewed in Gorbach, S.L., et al. [28]) and the like. Immunological therapy will also be  
30 effective in many cases to manage and alleviate symptoms caused by the disease agents described here. Antiviral agents include agents or compositions that directly bind to viral products and interfere with disease progress; and, excludes agents that do not  
35 impact directly on viral multiplication or viral titer. Antiviral agents do not include immunoregulatory agents that do not directly affect



these agents are preferentially phosphorylated by viral thymidine kinase (TK), if one is present, and/or have higher affinity for viral DNA polymerase than for the cellular DNA polymerases, resulting in selective antiviral activity. Where a nucleoside analogue is incorporated into the viral DNA, viral activity or reproduction may be affected in a variety of ways. For example, the analogue may act as a chain terminator, cause increased lability (e.g., susceptibility to breakage) of analogue-containing DNA, and/or impair the ability of the substituted DNA to act as template for transcription or replication (see, e.g., Balzarini et al. [11]).

It will be known to one of skill that, like many drugs, many of the agents useful for treatment of herpes virus infections are modified (*i.e.*, "activated") by the host, host cell, or virus-infected host cell metabolic enzymes. For example, acyclovir is triphosphorylated to its active form, with the first phosphorylation being carried out by the herpes virus thymidine kinase, when present. Other examples are the reported conversion of the compound HOE 602 to ganciclovir in a three-step metabolic pathway (Winkler et al. [95]) and the phosphorylation of ganciclovir to its active form by, e.g., a CMV nucleotide kinase. It will be apparent to one of skill that the specific metabolic capabilities of a virus can affect the sensitivity of that virus to specific drugs, and is one factor in the choice of an antiviral drug. The mechanism of action of certain anti-herpesvirus agents is discussed in De Clercq [22] and in other references cited *supra* and *infra*, all of which are incorporated by reference herein.

Anti-herpesvirus medications suitable for treating viral induced KS include, but are not limited to,

nucleoside analogs including acyclic nucleoside  
phosphonate analogs (e.g.,  
phosphonylethoxyalkylpurines and -pyrimidines), and  
cyclic nucleoside analogs. These include drugs such  
5 as: vidarabine (9- $\beta$ -D-arabinofuranosyladenine; adenine  
arabinoside, ara-A, Vira-A, Parke-Davis); 1- $\beta$ -D-  
arabinofuranosyluracil (ara-U); 1- $\beta$ -D-  
arabinofuranosyl-cytosine (ara-C); HPMPA [(S)-1-[3-  
10 hydroxy-2-(phosphonylmethoxy)propyl]cytosine (e.g., GS  
504 Gilead Science)] and its cyclic form (cHPMPA);  
HPMPA [(S)-9-(3-hydroxy-2-  
phosphonylmethoxypropyl)adenine] and its cyclic form  
(cHPMPA); (S)-HPMPDAP [(S)-9-(3-hydroxy-2-  
15 phosphonylmethoxypropyl)-2,6-diaminopurine]; PMEDAP  
[9-(2-phosphonyl-methoxyethyl)-2,6-diaminopurine]; HOE  
602 [2-amino-9-(1,3-bis(isopropoxy)-2-  
propoxymethyl)purine]; PMEPA [9-(2-  
phosphonylmethoxyethyl)adenine]; bromovinyl-  
20 deoxyuridine (Burns and Sandford. [21]); 1- $\beta$ -D-  
arabinofuranosyl-E-5-(2-bromovinyl)-uridine or -2'-  
deoxyuridine; BVaraU (1- $\beta$ -D-arabinofuranosyl-E-5-(2-  
bromovinyl)-uracil, brovavir, Bristol-Myers Squibb,  
Yamsa Shoyu); BVDU [(E)-5-(2-bromovinyl)-2'-  
25 deoxyuridine, brivudin, e.g., Helpin] and its  
carbocyclic analogue (in which the sugar moiety is  
replaced by a cyclopentane ring); IVDU [(E)-5-(2-  
iodovinyl)-2'-deoxyuridine] and its carbocyclic  
analogue, C-IVDU (Balzarini et al. [11]); and 5-  
30 mercutithio analogs of 2'-deoxyuridine (Holliday, J.,  
and Williams, M.V. [38]); acyclovir [9-([2-  
hydroxyethoxy)methyl]guanine; e.g., Zovirax (Burroughs  
Wellcome)]; penciclovir (9-[4-hydroxy-2-  
(hydroxymethyl)butyl]-guanine); ganciclovir [(9-[1,3-  
35 dihydroxy-2 propoxymethyl]-guanine) e.g., Cymevene,  
Cytovene (Syntex), DHPG (Stals et al. [89]);  
isopropylether derivatives of ganciclovir (see, e.g.,  
Winkelmann et al. [94]); cygalovir; famciclovir [2-

amino-9-(4-acetoxy-3-(acetoxymethyl)but-1-yl)purine  
(Smithkline Beecham)]; valacyclovir (Burroughs  
Wellcome); desciclovir [(2-amino-9-(2-  
ethoxymethyl)purine)] and 2-amino-9-(2-  
5 hydroxyethoxymethyl)-9H-purine, prodrugs of  
acyclovir]; CDG (carbocyclic 2'-deoxyguanosine); and  
purine nucleosides with the pentafuranosyl ring  
replaced by a cyclo butane ring (e.g., cyclobut-A [(+)  
)-9-[1 $\beta$ , 2 $\alpha$ , 3 $\beta$ ]-2,3-bis(hydroxymethyl)-1-  
10 cyclobutyl]adenine], cyclobut-G [(+)-9-[1 $\beta$ , 2 $\alpha$ , 3 $\beta$ ]-  
2,3-bis(hydroxymethyl)-1-cyclobutyl]guanine], BHCG  
[(R)-(1 $\alpha$ , 2 $\beta$ , 1 $\alpha$ )-9-(2,3-  
bis(hydroxymethyl)cyclobutyl]guanine], and an active  
isomer of racemic BHCG, SQ 34,514 [1R-1 $\alpha$ , 2 $\beta$ , 3 $\alpha$ )-2-  
15 amino-9-[2,3-bis(hydroxymethyl)cyclobutyl]-6H-purin-6-  
one (see, Braitman et al.(1991) [20]]. Certain of  
these antiherpesviral agents are discussed in Gorach  
et al. [28]; Saunders et al. [82]; Yamanaka et al.,  
[96]; Greenspan et al. [29], all of which are  
20 incorporated by reference herein.

Triciribine and triciribine monophosphate are potent  
inhibitors against herpes viruses. (Ickes et al. [43],  
incorporated by reference herein), HIV-1 and HIV-2  
25 (Kucera et al. [51], incorporated by reference herein)  
and are additional nucleoside analogs that may be used  
to treat KS. An exemplary protocol for these agents  
is an intravenous injection of about 0.35 mg/meter<sup>2</sup>  
(0.7 mg/kg) once weekly or every other week for at  
30 least two doses, preferably up to about four to eight  
weeks.

Acyclovir and ganciclovir are of interest because of  
their accepted use in clinical settings. Acyclovir,  
35 an acyclic analogue of guanine, is phosphorylated by  
a herpesvirus thymidine kinase and undergoes further  
phosphorylation to be incorporated as a chain

terminator by the viral DNA polymerase during viral replication. It has therapeutic activity against a broad range of herpesviruses, Herpes simplex Types 1 and 2, Varicella- Zoster, Cytomegalovirus, and Epstein-Barr Virus, and is used to treat disease such as herpes encephalitis, neonatal herpesvirus infections, chickenpox in immunocompromised hosts, herpes zoster recurrences, CMV retinitis, EBV infections, chronic fatigue syndrome, and hairy leukoplakia in AIDS patients. Exemplary intravenous dosages or oral dosages are 250 mg/kg/m<sup>2</sup> body surface area, every 8 hours for 7 days, or maintenance doses of 200-400 mg IV or orally twice a day to suppress recurrence. Ganciclovir has been shown to be more active than acyclovir against some herpesviruses. See, e.g., Oren and Soble [73]. Treatment protocols for ganciclovir are 5 mg/kg twice a day IV or 2.5 mg/kg three times a day for 10-14 days. Maintenance doses are 5-6 mg/kg for 5-7 days.

Also of interest is HPMPC. HPMPC is reported to be more active than either acyclovir or ganciclovir in the chemotherapy and prophylaxis of various HSV-1, HSV-2, TK- HSV, VZV or CMV infections in animal models ([22], *supra*).

Nucleoside analogs such as BVaraU are potent inhibitors of HSV-1, EBV, and VZV that have greater activity than acyclovir in animal models of encephalitis. FIAC (fluoridoarbinosyl cytosine) and its related fluoroethyl and iodo compounds (e.g., FEAU, FIAU) have potent selective activity against herpesviruses, and HPMPA ((S)-1-([3-hydroxy-2-phosphorylmethoxy]propyl)adenine) has been demonstrated to be more potent against HSV and CMV than acyclovir or ganciclovir and are of choice in advanced cases of KS. Cladribine (2-

chlorodeoxyadenosine) is another nucleoside analogue known as a highly specific antilymphocyte agent (i.e., a immunosuppressive drug).

5 Other useful antiviral agents include: 5-thien-2-yl-2'-deoxyuridine derivatives, e.g., BTDU [5-(5-bromothien-2-yl)-2'-deoxyuridine] and CTDU [5-(5-chlorothien-2-yl)-2'-deoxyuridine]; and OXT-A [9-(2-deoxy-2-hydroxymethyl- $\beta$ -D-erythro-oxetanosyl)adenine]  
 10 and OXT-G [9-(2-deoxy-2-hydroxymethyl- $\beta$ -D-erythro-oxetanosyl)guanine]. Although OXT-G is believed to act by inhibiting viral DNA synthesis its mechanism of action has not yet been elucidated. These and other compounds are described in Andrei et al. [5] which is  
 15 incorporated by reference herein. Additional antiviral purine derivatives useful in treating herpesvirus infections are disclosed in US Pat. 5,108,994 (assigned to Beecham Group P.L.C.). 6-Methoxypurine arabinoside (ara-M; Burroughs Wellcome)  
 20 is a potent inhibitor of varicella-zoster virus, and will be useful for treatment of KS.

Certain thymidine analogs [e.g., idoxuridine (5-ido-2'-deoxyuridine)] and triflurothymidine) have  
 25 antiherpes viral activity, but due to their systemic toxicity, are largely used for topical herpesviral infections, including HSV stromal keratitis and uveitis, and are not preferred here unless other options are ruled out.

30 Other useful antiviral agents that have demonstrated antiherpes viral activity include foscarnet sodium (trisodium phosphonoformate, PFA, Foscavir (Astra)) and phosphonoacetic acid (PAA). Foscarnet is an  
 35 inorganic pyrophosphate analogue that acts by competitively blocking the pyrophosphate-binding site of DNA polymerase. These agents which block DNA



polymerase directly without processing by viral thymidine kinase. Foscarnet is reported to be less toxic than PAA.

- 5                   ii) Agents that target viral proteins other than DNA polymerase or other viral functions.

10                   Although applicants do not intend to be bound by a particular mechanism of antiviral action, the antiherpes-virus agents described above are believed to act through inhibition of viral DNA polymerase. However, viral replication requires not only the replication of the viral nucleic acid but also the production of viral proteins and other essential components. Accordingly, the present invention contemplates treatment of KS by the inhibition of viral proliferation by targeting viral proteins other than DNA polymerase (e.g., by inhibition of their synthesis or activity, or destruction of viral proteins after their synthesis). For example, administration of agents that inhibit a viral serine protease, e.g., such as one important in development of the viral capsid will be useful in treatment of viral induced KS.

15

20

25

Other viral enzyme targets include: OMP decarboxylase inhibitors (a target of, e.g., parazofurin), CTP synthetase inhibitors (targets of, e.g., cyclopentenylcytosine), IMP dehydrogenase, ribonucleotide reductase (a target of, e.g., carboxyl-containing N-alkyldipeptides as described in U.S. Patent No. 5,110,799 (Tolman et al., Merck)), thymidine kinase (a target of, e.g., 1-[2-(hydroxymethyl)cycloalkylmethyl]-5-substituted -uracils and -guanines as described in, e.g., U.S. Patent Nos. 4,863,927 and 4,782,062 (Tolman et al.;

30

35

Merck)) as well as other enzymes. It will be apparent to one of ordinary skill in the art that there are additional viral proteins, both characterized and as yet to be discovered, that can serve as target for antiviral agents.

iv) Other agents and modes of antiviral action.

10 Kutapressin is a liver derivative available from Schwarz Parma of Milwaukee, Wisconsin in an injectable form of 25 mg/ml. The recommended dosage for herpesviruses is from 200 to 25 mg/ml per day for an average adult of 150 pounds.

15 Poly(I)·Poly(C<sub>12</sub>U), an accepted antiviral drug known as Ampligen from HEM Pharmaceuticals of Rockville, MD has been shown to inhibit herpesviruses and is another antiviral agent suitable for treating KS. Intravenous injection is the preferred route of administration. 20 Dosages from about 100 to 600 mg/m<sup>2</sup> are administered two to three times weekly to adults averaging 150 pounds. It is best to administer at least 200 mg/m<sup>2</sup> per week.

25 Other antiviral agents reported to show activity against herpes viruses (e.g., varicella zoster and herpes simplex) and will be useful for the treatment of herpesvirus-induced KS include mappicine ketone 30 (SmithKline Beecham); Compounds A,79296 and A,73209 (Abbott) for varicella zoster, and Compound 882C87 (Burroughs Wellcome) [see, *The Pink Sheet* 55(20) May 17, 1993].

35 Interferon is known inhibit replication of herpes viruses. See [73], *supra*. Interferon has known toxicity problems and it is expected that second

005230-02E960

generation derivatives will soon be available that will retain interferon's antiviral properties but have reduced side effects.

5 It is also contemplated that herpes virus-induced KS may be treated by administering a herpesvirus reactivating agent to induce reactivation of the latent virus. Preferably the reactivation is combined with simultaneous or sequential administration of an anti-herpesvirus agent. Controlled reactivation over 10 a short period of time or reactivation in the presence of an antiviral agent is believed to minimize the adverse effects of certain herpesvirus infections (e.g., as discussed in PCT Application WO 93/04683). 15 Reactivating agents include agents such as estrogen, phorbol esters, forskolin and  $\beta$ -adrenergic blocking agents.

Agents useful for treatment of herpesvirus infections and for treatment of herpesvirus-induced KS are 20 described in numerous U.S. Patents. For example, ganciclovir is an example of a antiviral guanine acyclic nucleotide of the type described in US Patent Nos. 4,355,032 and 4,603,219.

25 Acyclovir is an example of a class of antiviral purine derivatives, including 9 - (2 - hydroxyethylmethyl)adenine, of the type described in U.S. Pat. Nos. 4,287,188, 4,294,831 and 4,199,574.

30 Brivudin is an example of an antiviral deoxyuridine derivative of the type described in US Patent No. 4,424,211.

35 Vidarabine is an example of an antiviral purine nucleoside of the type described in British Pat. 1,159,290.

006250"0220360

Brovavir is an example of an antiviral deoxyuridine derivative of the type described in US Patent Nos. 4,542,210 and 4,386,076.

5 BHCG is an example of an antiviral carbocyclic nucleoside analogue of the type described in US Patent Nos. 5,153,352, 5,034,394 and 5,126,345.

10 HPMPC is an example of an antiviral phosphonyl methoxyalkyl derivative with of the type described in US Patent No. 5,142,051.

15 CDG (Carbocyclic 2'-deoxyguanosine) is an example of an antiviral carbocyclic nucleoside analogue of the type described in US Patent Nos. 4,543,255, 4,855,466, and 4,894,458.

Foscarnet is described in US Patent No. 4,339,445.

20 Trifluridine and its corresponding ribonucleoside is described in US Patent No. 3,201,387.

25 U.S. Patent No. 5,321,030 (Kaddurah-Daouk et al.; Amira) describes the use of creatine analogs as antiherpes viral agents. U.S. Patent No. 5,306,722 (Kim et al.; Bristol-Meyers Squibb) describes thymidine kinase inhibitors useful for treating HSV infections and for inhibiting herpes thymidine kinase. Other antiherpesvirus compositions are described in  
30 U.S. Patent Nos. 5,286,649 and 5,098,708 (Konishi et al., Bristol-Meyers Squibb) and 5,175,165 (Blumenkopf et al.; Burroughs Wellcome). U.S. Patent No. 4,880,820 (Ashton et al.; Merck) describes the  
35 antiherpes virus agent (S)-9-(2,3-dihydroxy-1-propoxymethyl)guanine.

006290-64740560

U.S. Patent No. 4,708,935 (Suhadolnik et al.; Research Corporation) describes a 3'-deoxyadenosine compound effective in inhibiting HSV and EBV. U.S. Patent No. 4,386,076 (Machida et al.; Yamasa Shoyu Kabushiki K a i s h a ) d e s c r i b e s u s e o f (E)-5-(2-halogenovinyl)-arabinofuranosyluracil as an antiherpesvirus agent. U.S. Patent No. 4,340,599 (Lieb et al.; Bayer Aktiengesellschaft) describes phosphonohydroxyacetic acid derivatives useful as antiherpes agents. U.S. Patent Nos. 4,093,715 and 4,093,716 (Lin et al. Research Corporation) describe 5'-amino-5'-deoxythymidine and 5-iodo-5'-amino-2',5'-dideoxycytidine as potent inhibitors of herpes simplex virus. U.S. Patent No. 4,069,382 (Baker et al.; Parke, Davis & Company) describes 9-(5-O-Acyl-beta-D-arabinofuranosyl)adenine compounds useful as antiviral agents. U.S. Patent No. 3,927,216 (Witkowski et al.) describes the use of 1,2,4-triazole-3-carboxamide and 1,2,4-triazole-3-thiocarboxamide for inhibiting herpes virus infections. Patent No. 5,179,093 (Afonso et al., Schering) describes quinoline-2,4-dione derivatives active against herpes simplex virus 1 and 2, cytomegalovirus and Epstein Barr virus.

25

## v) Inhibitory nucleic acid therapeutics

Also contemplated here are inhibitory nucleic acid therapeutics which can inhibit the activity of herpesviruses in patients with KS. Inhibitory nucleic acids may be single-stranded nucleic acids, which can specifically bind to a complementary nucleic acid sequence. By binding to the appropriate target sequence, an RNA-RNA, a DNA-DNA, or RNA-DNA duplex or triplex is formed. These nucleic acids are often termed "antisense" because they are usually complementary to the sense or coding strand of the

30

35

006250-62720360

gene, although recently approaches for use of "sense" nucleic acids have also been developed. The term "inhibitory nucleic acids" as used herein, refers to both "sense" and "antisense" nucleic acids.

5

By binding to the target nucleic acid, the inhibitory nucleic acid can inhibit the function of the target nucleic acid. This could, for example, be a result of blocking DNA transcription, processing or poly(A) addition to mRNA, DNA replication, translation, or promoting inhibitory mechanisms of the cells, such as promoting RNA degradation. Inhibitory nucleic acid methods therefore encompass a number of different approaches to altering expression of herpesvirus genes. These different types of inhibitory nucleic acid technology are described in Helene, C. and Toulme, J. [34], which is hereby incorporated by reference and is referred to hereinafter as "Helene and Toulme."

10  
15  
20

In brief, inhibitory nucleic acid therapy approaches can be classified into those that target DNA sequences, those that target RNA sequences (including pre-mRNA and mRNA), those that target proteins (sense strand approaches), and those that cause cleavage or chemical modification of the target nucleic acids.

25

Approaches targeting DNA fall into several categories. Nucleic acids can be designed to bind to the major groove of the duplex DNA to form a triple helical or "triplex" structure. Alternatively, inhibitory nucleic acids are designed to bind to regions of single stranded DNA resulting from the opening of the duplex DNA during replication or transcription. See Helene and Toulme.

30  
35

More commonly, inhibitory nucleic acids are designed to bind to mRNA or mRNA precursors. Inhibitory nucleic acids are used to prevent maturation of pre-mRNA. Inhibitory nucleic acids may be designed to  
5 interfere with RNA processing, splicing or translation.

The inhibitory nucleic acids can be targeted to mRNA. In this approach, the inhibitory nucleic acids are  
10 designed to specifically block translation of the encoded protein. Using this approach, the inhibitory nucleic acid can be used to selectively suppress certain cellular functions by inhibition of translation of mRNA encoding critical proteins. For  
15 example, an inhibitory nucleic acid complementary to regions of c-myc mRNA inhibits c-myc protein expression in a human promyelocytic leukemia cell line, HL60, which overexpresses the c-myc proto-oncogene. See Wickstrom E.L., et al. [93] and  
20 Harel-Bellan, A., et al. [31A]. As described in Helene and Toulme, inhibitory nucleic acids targeting mRNA have been shown to work by several different mechanisms to inhibit translation of the encoded protein(s).

25 The inhibitory nucleic acids introduced into the cell can also encompass the "sense" strand of the gene or mRNA to trap or compete for the enzymes or binding proteins involved in mRNA translation. See Helene and  
30 Toulme.

Lastly, the inhibitory nucleic acids can be used to induce chemical inactivation or cleavage of the target genes or mRNA. Chemical inactivation can occur by the  
35 induction of crosslinks between the inhibitory nucleic acid and the target nucleic acid within the cell. Other chemical modifications of the target nucleic

acids induced by appropriately derivatized inhibitory nucleic acids may also be used.

5 Cleavage, and therefore inactivation, of the target nucleic acids may be effected by attaching a substituent to the inhibitory nucleic acid which can be activated to induce cleavage reactions. The substituent can be one that affects either chemical, or enzymatic cleavage. Alternatively, cleavage can  
10 be induced by the use of ribozymes or catalytic RNA. In this approach, the inhibitory nucleic acids would comprise either naturally occurring RNA (ribozymes) or synthetic nucleic acids with catalytic activity.

15 The targeting of inhibitory nucleic acids to specific cells of the immune system by conjugation with targeting moieties binding receptors on the surface of these cells can be used for all of the above forms of inhibitory nucleic acid therapy. This invention  
20 encompasses all of the forms of inhibitory nucleic acid therapy as described above and as described in Helene and Toulme.

This invention relates to the targeting of inhibitory  
25 nucleic acids to sequences the human herpesvirus of the invention for use in treating KS. An example of an antiherpes virus inhibitory nucleic acid is ISIS 2922 (ISIS Pharmaceuticals) which has activity against CMV [see, *Biotechnology News* 14(14) p. 5].

30 A problem associated with inhibitory nucleic acid therapy is the effective delivery of the inhibitory nucleic acid to the target cell in vivo and the subsequent internalization of the inhibitory nucleic  
35 acid by that cell. This can be accomplished by linking the inhibitory nucleic acid to a targeting moiety to form a conjugate that binds to a specific



receptor on the surface of the target infected cell, and which is internalized after binding.

iii) Administration

5

The subjects to be treated or whose tissue may be used herein may be a mammal, or more specifically a human, horse, pig, rabbit, dog, monkey, or rodent. In the preferred embodiment the subject is a human.

10

The compositions are administered in a manner compatible with the dosage formulation, and in a therapeutically effective amount. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and are peculiar to each subject.

15

Suitable regimes for initial administration and booster shots are also variable, but are typified by an initial administration followed by repeated doses at one or more hour intervals by a subsequent injection or other administration.

20

As used herein administration means a method of administering to a subject. Such methods are well known to those skilled in the art and include, but are not limited to, administration topically, parenterally, orally, intravenously, intramuscularly, subcutaneously or by aerosol. Administration of the agent may be effected continuously or intermittently such that the therapeutic agent in the patient is effective to treat a subject with Kaposi's sarcoma or a subject infected with a DNA virus associated with Kaposi's sarcoma.

25

30

35

The antiviral compositions for treating herpesvirus-induced KS are preferably administered to human

patients via oral, intravenous or parenteral  
administrations and other systemic forms. Those of  
skill in the art will understand appropriate  
administration protocol for the individual  
5 compositions to be employed by the physician.

The pharmaceutical formulations or compositions of  
this invention may be in the dosage form of solid,  
semi-solid, or liquid such as, e.g., suspensions,  
10 aerosols or the like. Preferably the compositions are  
administered in unit dosage forms suitable for single  
administration of precise dosage amounts. The  
compositions may also include, depending on the  
formulation desired, pharmaceutically-acceptable, non-  
15 toxic carriers or diluents, which are defined as  
vehicles commonly used to formulate pharmaceutical  
compositions for animal or human administration. The  
diluent is selected so as not to affect the biological  
activity of the combination. Examples of such  
20 diluents are distilled water, physiological saline,  
Ringer's solution, dextrose solution, and Hank's  
solution. In addition, the pharmaceutical composition  
or formulation may also include other carriers,  
adjuvants; or nontoxic, nontherapeutic, nonimmunogenic  
25 stabilizers and the like. Effective amounts of such  
diluent or carrier are those amounts which are  
effective to obtain a pharmaceutically acceptable  
formulation in terms of solubility of components, or  
biological activity, etc.

30

#### V. Immunological Approaches to Therapy.

Having identified a primary causal agent of KS in  
humans as a novel human herpesvirus, there are  
35 immunosuppressive therapies that can modulate the  
immunologic dysfunction that arises from the presence  
of viral infected tissue. In particular, agents that

block the immunological attack of the viral infected cells will ameliorate the symptoms of KS and/or reduce the disease progress. Such therapies include antibodies that specifically block the targeting of viral infected cells. Such agents include antibodies which bind to cytokines that upregulate the immune system to target viral infected cells.

The antibody may be administered to a patient either singly or in a cocktail containing two or more antibodies, other therapeutic agents, compositions, or the like, including, but not limited to, immunosuppressive agents, potentiators and side-effect relieving agents. Of particular interest are immunosuppressive agents useful in suppressing allergic reactions of a host. Immunosuppressive agents of interest include prednisone, prednisolone, DECADRON (Merck, Sharp & Dohme, West Point, PA), cyclophosphamide, cyclosporine, 6-mercaptopurine, methotrexate, azathioprine and i.v. gamma globulin or their combination. Potentiators of interest include monensin, ammonium chloride and chloroquine. All of these agents are administered in generally accepted efficacious dose ranges such as those disclosed in the *Physician Desk Reference*, 41st Ed. (1987), Publisher Edward R. Barnhart, New Jersey.

Immune globulin from persons previously infected with human herpesviruses or related viruses can be obtained using standard techniques. Appropriate titers of antibodies are known for this therapy and are readily applied to the treatment of KS. Immune globulin can be administered via parenteral injection or by intrathecal shunt. In brief, immune globulin preparations may be obtained from individual donors who are screened for antibodies to the KS-associated human herpesvirus, and plasmas from high-titered

donors are pooled. Alternatively, plasmas from donors are pooled and then tested for antibodies to the human herpesvirus of the invention; high-titered pools are then selected for use in KS patients.

5

Antibodies may be formulated into an injectable preparation. Parenteral formulations are known and are suitable for use in the invention, preferably for i.m. or i.v. administration. The formulations containing therapeutically effective amounts of antibodies or immunotoxins are either sterile liquid solutions, liquid suspensions or lyophilized versions and optionally contain stabilizers or excipients. Lyophilized compositions are reconstituted with suitable diluents, e.g., water for injection, saline, 0.3% glycine and the like, at a level of about from .01 mg/kg of host body weight to 10 mg/kg where appropriate. Typically, the pharmaceutical compositions containing the antibodies or immunotoxins will be administered in a therapeutically effective dose in a range of from about .01 mg/kg to about 5 mg/kg of the treated mammal. A preferred therapeutically effective dose of the pharmaceutical composition containing antibody or immunotoxin will be in a range of from about 0.01 mg/kg to about 0.5 mg/kg body weight of the treated mammal administered over several days to two weeks by daily intravenous infusion, each given over a one hour period, in a sequential patient dose-escalation regimen.

30

Antibody may be administered systemically by injection i.m., subcutaneously or intraperitoneally or directly into KS lesions. The dose will be dependent upon the properties of the antibody or immunotoxin employed, e.g., its activity and biological half-life, the concentration of antibody in the formulation, the site and rate of dosage, the clinical tolerance of the

35

DDB-2013-04-23-10:50:50

patient involved, the disease afflicting the patient and the like as is well within the skill of the physician.

5 The antibody of the present invention may be administered in solution. The pH of the solution should be in the range of pH 5 to 9.5, preferably pH 6.5 to 7.5. The antibody or derivatives thereof should be in a solution having a suitable  
10 pharmaceutically acceptable buffer such as phosphate, tris (hydroxymethyl) aminomethane-HCl or citrate and the like. Buffer concentrations should be in the range of 1 to 100 mM. The solution of antibody may also contain a salt, such as sodium chloride or  
15 potassium chloride in a concentration of 50 to 150 mM. An effective amount of a stabilizing agent such as an albumin, a globulin, a gelatin, a protamine or a salt of protamine may also be included and may be added to a solution containing antibody or immunotoxin or to  
20 the composition from which the solution is prepared.

Systemic administration of antibody is made daily, generally by intramuscular injection, although intravascular infusion is acceptable. Administration  
25 may also be intranasal or by other nonparenteral routes. Antibody or immunotoxin may also be administered via microspheres, liposomes or other microparticulate delivery systems placed in certain tissues including blood.

30 In therapeutic applications, the dosages of compounds used in accordance with the invention vary depending on the class of compound and the condition being treated. The age, weight, and clinical condition of  
35 the recipient patient; and the experience and judgment of the clinician or practitioner administering the therapy are among the factors affecting the selected

dosage. For example, the dosage of an immunoglobulin can range from about 0.1 milligram per kilogram of body weight per day to about 10 mg/kg per day for polyclonal antibodies and about 5% to about 20% of that amount for monoclonal antibodies. In such a case, the immunoglobulin can be administered once daily as an intravenous infusion. Preferably, the dosage is repeated daily until either a therapeutic result is achieved or until side effects warrant discontinuation of therapy. Generally, the dose should be sufficient to treat or ameliorate symptoms or signs of KS without producing unacceptable toxicity to the patient.

An effective amount of the compound is that which provides either subjective relief of a symptom(s) or an objectively identifiable improvement as noted by the clinician or other qualified observer. The dosing range varies with the compound used, the route of administration and the potency of the particular compound.

#### VI. Vaccines and Prophylaxis for KS

This invention provides a method of vaccinating a subject against Kaposi's sarcoma, comprising administering to the subject an effective amount of the peptide or polypeptide encoded by the isolated DNA molecule, and a suitable acceptable carrier, thereby vaccinating the subject. In one embodiment naked DNA is administering to the subject in an effective amount to vaccinate a subject against Kaposi's sarcoma.

This invention provides a method of immunizing a subject against a disease caused by the DNA herpesvirus associated with Kaposi's sarcoma which

comprises administering to the subject an effective immunizing dose of the isolated herpesvirus vaccine.

#### A. Vaccines

5

The invention also provides substances suitable for use as vaccines for the prevention of KS and methods for administering them. The vaccines are directed against the human herpesvirus of the invention, and most preferably comprise antigen obtained from the KS-associated human herpesvirus.

Vaccines can be made recombinantly. Typically, a vaccine will include from about 1 to about 50 micrograms of antigen or antigenic protein or peptide. More preferably, the amount of protein is from about 15 to about 45 micrograms. Typically, the vaccine is formulated so that a dose includes about 0.5 milliliters. The vaccine may be administered by any route known in the art. Preferably, the route is parenteral. More preferably, it is subcutaneous or intramuscular.

There are a number of strategies for amplifying an antigen's effectiveness, particularly as related to the art of vaccines. For example, cyclization or circularization of a peptide can increase the peptide's antigenic and immunogenic potency. See U.S. Pat. No. 5,001,049 which is incorporated by reference herein. More conventionally, an antigen can be conjugated to a suitable carrier, usually a protein molecule. This procedure has several facets. It can allow multiple copies of an antigen, such as a peptide, to be conjugated to a single larger carrier molecule. Additionally, the carrier may possess properties which facilitate transport, binding, absorption or transfer of the antigen.

For parenteral administration, such as subcutaneous injection, examples of suitable carriers are the tetanus toxoid, the diphtheria toxoid, serum albumin and lamprey, or keyhole limpet, hemocyanin because they provide the resultant conjugate with minimum genetic restriction. Conjugates including these universal carriers can function as T cell clone activators in individuals having very different gene sets.

10

The conjugation between a peptide and a carrier can be accomplished using one of the methods known in the art. Specifically, the conjugation can use bifunctional cross-linkers as binding agents as detailed, for example, by Means and Feeney, "A recent review of protein modification techniques," *Bioconjugate Chem.* 1:2-12 (1990).

15

Vaccines against a number of the Herpesviruses have been successfully developed. Vaccines against Varicella-Zoster Virus using a live attenuated Oka strain is effective in preventing herpes zoster in the elderly, and in preventing chickenpox in both immunocompromised and normal children (Hardy, I., et al. [30]; Hardy, I. et al. [31]; Levin, M.J. et al. [54]; Gershon, A.A. [26]). Vaccines against Herpes simplex Types 1 and 2 are also commercially available with some success in protection against primary disease, but have been less successful in preventing the establishment of latent infection in sensory ganglia (Roizman, B. [78]; Skinner, G.R. et al. [87]).

20

25

30

Vaccines against the human herpesvirus can be made by isolating extracellular viral particles from infected cell cultures, inactivating the virus with formaldehyde followed by ultracentrifugation to concentrate the viral particles and remove the

35

ORIGINAL PAGE



formaldehyde, and immunizing individuals with 2 or 3 doses containing  $1 \times 10^9$  virus particles (Skinner, G.R. et al. [86]). Alternatively, envelope glycoproteins can be expressed in E. coli or transfected into stable mammalian cell lines, the proteins can be purified and used for vaccination (Lasky, L.A. [53]). MHC-binding peptides from cells infected with the human herpesvirus can be identified for vaccine candidates per the methodology of [61], supra.

The antigen may be combined or mixed with various solutions and other compounds as is known in the art. For example, it may be administered in water, saline or buffered vehicles with or without various adjuvants or immunodiluting agents. Examples of such adjuvants or agents include aluminum hydroxide, aluminum phosphate, aluminum potassium sulfate (alum), beryllium sulfate, silica, kaolin, carbon, water-in-oil emulsions, oil-in-water emulsions, muramyl dipeptide, bacterial endotoxin, lipid X, *Corynebacterium parvum* (*Propionibacterium acnes*), *Bordetella pertussis*, polyribonucleotides, sodium alginate, lanolin, lysolecithin, vitamin A, saponin, liposomes, levamisole, DEAE-dextran, blocked copolymers or other synthetic adjuvants. Such adjuvants are available commercially from various sources, for example, Merck Adjuvant 65 (Merck and Company, Inc., Rahway, N.J.) or Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, Michigan). Other suitable adjuvants are Amphigen (oil-in-water), Alhydrogel (aluminum hydroxide), or a mixture of Amphigen and Alhydrogel. Only aluminum is approved for human use.

The proportion of antigen and adjuvant can be varied over a broad range so long as both are present in effective amounts. For example, aluminum hydroxide

can be present in an amount of about 0.5% of the vaccine mixture (Al<sub>2</sub>O<sub>3</sub> basis). On a per-dose basis, the amount of the antigen can range from about 0.1 μg to about 100 μg protein per patient. A preferable range is from about 1 μg to about 50 μg per dose. A more preferred range is about 15 μg to about 45 μg. A suitable dose size is about 0.5 ml. Accordingly, a dose for intramuscular injection, for example, would comprise 0.5 ml containing 45 μg of antigen in admixture with 0.5% aluminum hydroxide. After formulation, the vaccine may be incorporated into a sterile container which is then sealed and stored at a low temperature, for example 4°C, or it may be freeze-dried. Lyophilization permits long-term storage in a stabilized form.

The vaccines may be administered by any conventional method for the administration of vaccines including oral and parenteral (e.g., subcutaneous or intramuscular) injection. Intramuscular administration is preferred. The treatment may consist of a single dose of vaccine or a plurality of doses over a period of time. It is preferred that the dose be given to a human patient within the first 8 months of life. The antigen of the invention can be combined with appropriate doses of compounds including influenza antigens, such as influenza type A antigens. Also, the antigen could be a component of a recombinant vaccine which could be adaptable for oral administration.

Vaccines of the invention may be combined with other vaccines for other diseases to produce multivalent vaccines. A pharmaceutically effective amount of the antigen can be employed with a pharmaceutically acceptable carrier such as a protein or diluent useful for the vaccination of mammals, particularly humans.

Other vaccines may be prepared according to methods well-known to those skilled in the art.

Those of skill will readily recognize that it is only  
5 necessary to expose a mammal to appropriate epitopes  
in order to elicit effective immunoprotection. The  
epitopes are typically segments of amino acids which  
are a small portion of the whole protein. Using  
10 recombinant genetics, it is routine to alter a natural  
protein's primary structure to create derivatives  
embracing epitopes that are identical to or  
substantially the same as (immunologically equivalent  
to) the naturally occurring epitopes. Such  
15 derivatives may include peptide fragments, amino acid  
substitutions, amino acid deletions and amino acid  
additions of the amino acid sequence for the viral  
proteins from the human herpesvirus. For example, it  
is known in the protein art that certain amino acid  
20 residues can be substituted with amino acids of  
similar size and polarity without an undue effect upon  
the biological activity of the protein. The human  
herpesvirus proteins have significant tertiary  
structure and the epitopes are usually conformational.  
Thus, modifications should generally preserve  
25 conformation to produce a protective immune response.

#### B. Antibody Prophylaxis

Therapeutic, intravenous, polyclonal or monoclonal  
30 antibodies can be used as a mode of passive  
immunotherapy of herpesviral diseases including  
perinatal varicella and CMV. Immune globulin from  
persons previously infected with the human herpesvirus  
and bearing a suitably high titer of antibodies  
35 against the virus can be given in combination with  
antiviral agents (e.g. ganciclovir), or in combination  
with other modes of immunotherapy that are currently

being evaluated for the treatment of KS, which are targeted to modulating the immune response (i.e. treatment with copolymer-1, antiidiotypic monoclonal antibodies, T cell "vaccination"). Antibodies to human herpesvirus can be administered to the patient as described herein. Antibodies specific for an epitope expressed on cells infected with the human herpesvirus are preferred and can be obtained as described above.

A polypeptide, analog or active fragment can be formulated into the therapeutic composition as neutralized pharmaceutically acceptable salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide or antibody molecule) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed from the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

### C. Monitoring therapeutic efficacy

This invention provides a method for monitoring the therapeutic efficacy of treatment for Kaposi's sarcoma, which comprises determining in a first sample from a subject with Kaposi's sarcoma the presence of the isolated DNA molecule, administering to the subject a therapeutic amount of an agent such that the agent is contacted to the cell in a sample, determining after a suitable period of time the amount of the isolated DNA molecule in the second sample from

being evaluated for the treatment of KS, which are targeted to modulating the immune response (i.e. treatment with copolymer-1, antiidiotypic monoclonal antibodies, T cell "vaccination"). Antibodies to human herpesvirus can be administered to the patient as described herein. Antibodies specific for an epitope expressed on cells infected with the human herpesvirus are preferred and can be obtained as described above.

10

A polypeptide, analog or active fragment can be formulated into the therapeutic composition as neutralized pharmaceutically acceptable salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide or antibody molecule) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed from the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

20  
25

### C. Monitoring therapeutic efficacy

This invention provides a method for monitoring the therapeutic efficacy of treatment for Kaposi's sarcoma, which comprises determining in a first sample from a subject with Kaposi's sarcoma the presence of the isolated DNA molecule, administering to the subject a therapeutic amount of an agent such that the agent is contacted to the cell in a sample, determining after a suitable period of time the amount of the isolated DNA molecule in the second sample from

30  
35

006290" 024050

the treated subject, and comparing the amount of isolated DNA molecule determined in the first sample with the amount determined in the second sample, a difference indicating the effectiveness of the agent, thereby monitoring the therapeutic efficacy of treatment for Kaposi's sarcoma. As defined herein "amount" is viral load or copy number. Methods of determining viral load or copy number are known to those skilled in the art.

VII. Screening Assays For Pharmaceutical Agents of Interest in Alleviating the Symptoms of KS.

Since an agent involved in the causation or progression of KS has been identified and described here, assays directed to identifying potential pharmaceutical agents that inhibit the biological activity of the agent are possible. KS drug screening assays which determine whether or not a drug has activity against the virus described herein are contemplated in this invention. Such assays comprise incubating a compound to be evaluated for use in KS treatment with cells which express the KS associated human herpesvirus proteins or peptides and determining therefrom the effect of the compound on the activity of such agent. In vitro assays in which the virus is maintained in suitable cell culture are preferred, though in vivo animal models would also be effective.

Compounds with activity against the agent of interest or peptides from such agent can be screened in in vitro as well as in vivo assay systems. In vitro assays include infecting peripheral blood leukocytes or susceptible T cell lines such as MT-4 with the agent of interest in the presence of varying concentrations of compounds targeted against viral replication, including nucleoside analogs, chain

terminators, antisense oligonucleotides and random polypeptides (Asada, H. et al. [7]; Kikuta et al. [48] both incorporated by reference herein). Infected cultures and their supernatants can be assayed for the total amount of virus including the presence of the viral genome by quantitative PCR, by dot blot assays, or by using immunologic methods. For example, a culture of susceptible cells could be infected with the human herpesvirus in the presence of various concentrations of drug, fixed on slides after a period of days, and examined for viral antigen by indirect immunofluorescence with monoclonal antibodies to viral peptides ([48], supra. Alternatively, chemically adhered MT-4 cell monolayers can be used for an infectious agent assay using indirect immunofluorescent antibody staining to search for focus reduction (Higashi, K. et al. [36], incorporated by reference herein).

As an alternative to whole cell in vitro assays, purified enzymes isolated from the human herpesvirus can be used as targets for rational drug design to determine the effect of the potential drug on enzyme activity, such as thymidine phosphotransferase or DNA polymerase. The genes for these two enzymes are provided herein. A measure of enzyme activity indicates effect on the agent itself.

Drug screens using herpes viral products are known and have been previously described in EP 0514830 (herpes proteases) and WO 94/04920 (U<sub>13</sub> gene product).

This invention provides an assay for screening anti-KS chemotherapeutics. Infected cells can be incubated in the presence of a chemical agent that is a potential chemotherapeutic against KS (e.g. acyclo-guanosine). The level of virus in the cells is then determined

5 after several days by IFA for antigens or Southern blotting for viral genome or Northern blotting for MRNA and compared to control cells. This assay can quickly screen large numbers of chemical compounds that may be useful against KS.

10 Further, this invention provides an assay system that is employed to identify drugs or other molecules capable of binding to the DNA molecule or proteins, either in the cytoplasm or in the nucleus, thereby inhibiting or potentiating transcriptional activity. Such assay would be useful in the development of drugs that would be specific against particular cellular activity, or that would potentiate such activity, in  
15 time or in level of activity.

20 This invention is further illustrated in the Experimental Details section which follows. This section is set forth to aid in an understanding of the invention but is not intended to, and should not be construed to, limit in any way the invention as set forth in the claims which follow thereafter.

25 **EXPERIMENTAL DETAILS SECTION I:**

**Experiment 1: Representational difference analysis (RDA) to identify and characterize unique DNA sequences in KS tissue**

30 To search for foreign DNA sequences belonging to an infectious agent in AIDS-KS, representational difference analysis (RDA) was employed to identify and characterize unique DNA sequences in KS tissue that are either absent or present in low copy number in  
35 non-diseased tissue obtained from the same patient [58]. This method can detect adenovirus genome added in single copy to human DNA but has not been used to



identify previously uncultured infectious agents. RDA is performed by making simplified "representations" of genomes from diseased and normal tissues from the same individual through PCR amplification of short restriction fragments. The DNA representation from the diseased tissue is then ligated to a priming sequence and hybridized to an excess of unligated, normal tissue DNA representation. Only unique sequences found in the diseased tissue have priming sequences on both DNA strands and are preferentially amplified during subsequent rounds of PCR amplification. This process can be repeated using different ligated priming sequences to enrich the sample for unique DNA sequences that are only found in the tissue of interest.

DNA (10  $\mu$ g) extracted from both the KS lesion and unaffected tissue were separately digested to completion with Bam HI (20 units/ $\mu$ g) at 37° C for 2 hours and 2  $\mu$ g of digestion fragments were ligated to NBam12 and NBam24 priming sequences [primer sequences described in 58]. Thirty cycles of PCR amplification were performed to amplify "representations" of both genomes. After construction of the genomic representations, KS tester amplicons between 150 and 1500 bp were isolated from an agarose gel and NBam priming sequences were removed by digestion with Bam HI. To search for unique DNA sequences not found in non-KS driver DNA, a second set of priming sequences (JBam12 and JBam24) was ligated onto only the KS tester DNA amplicons (Figure 1, lane 1). 0.2  $\mu$ g of ligated KS lesion amplicons were hybridized to 20  $\mu$ g of unligated, normal tissue representational amplicons. An aliquot of the hybridization product was then subjected to 10 cycles of PCR amplification using JBam24, followed by mung bean nuclease digestion. An aliquot of the mung bean-treated

difference product was then subjected to 15 more  
cycles of PCR with the JBam24 primer (Figure 1, lane  
2). Amplification products were redigested with Bam  
HI and 200 ng of the digested product was ligated to  
5 RBam12 and RBam24 primer sets for a second round of  
hybridization and PCR amplification (Figure 1, lane  
3). This enrichment procedure was repeated a third  
time using the JBam primer set (Figure 1, lane 4).  
Both the original driver and the tester DNA samples  
10 (Table 2, Patient A) were subsequently found to  
contain the AIDS-KS specific sequences KS330Bam and  
KS631Bam (previously identified as KS627Bam)  
indicating that RDA can be successfully employed when  
the target sequences are present in unequal copy  
15 number in both tissues.

The initial round of DNA amplification-hybridization  
from KS and normal tissue resulted in a diffuse  
banding pattern (Figure 1, lane 2), but four bands at  
20 approximately 380, 450, 540 and 680 bp were  
identifiable after the second amplification-  
hybridization (Figure 1, lane 3). These bands became  
discrete after a third round of amplification-  
hybridization (Figure 1, lane 4). Control RDA,  
25 performed by hybridizing DNA extracted from AIDS-KS  
tissue against itself, produced a single band at  
approximately 540 bp (Figure 1, lane 5). The four KS-  
associated bands (designated KS330Bam, KS390Bam,  
KS480Bam, KS627Bam after digestion of the two flanking  
30 28 bp ligated priming sequences with Bam HI) were gel  
purified and cloned by insertion into the pCRII  
vector. PCR products were cloned in the pCRII vector  
using the TA cloning system (Invitrogen Corporation,  
San Diego, CA).

**Experiment 2: Determination of the specificity of  
AIDS-KS unique sequences.**

5 To determine the specificity of these sequences for  
AIDS-KS, random-primed <sup>32</sup>P-labeled inserts were  
hybridized to Southern blots of DNA extracted from  
cryopreserved tissues obtained from patients with and  
without AIDS. All AIDS-KS specimens were examined  
microscopically for morphologic confirmation of KS and  
10 immunohistochemically for Factor VIII, Ulex europaeus  
and CD34 antigen expression. One of the AIDS-KS  
specimens was apparently mislabeled since KS tissue  
was not detected on microscopic examination but was  
included in the KS specimen group for purposes of  
15 statistical analysis. Control tissues used for  
comparison to the KS lesions included 56 lymphomas  
from patients with and without AIDS, 19 hyperplastic  
lymph nodes from patients with and without AIDS, 5  
vascular tumors from nonAIDS patients and 13 tissues  
20 infected with opportunistic infections that commonly  
occur in AIDS patients. Control DNA was also  
extracted from a consecutive series of 49 surgical  
biopsy specimens from patients without AIDS.  
Additional clinical and demographic information on the  
25 specimens was not collected to preserve patient  
confidentiality.

The tissues, listed in Table 1, were collected from  
diagnostic biopsies and autopsies between 1983 and  
30 1993 and stored at -70°C. Each tissue sample was from  
a different patient, except as noted in Table 1. Most  
of the 27 KS specimens were from lymph nodes dissected  
under surgical conditions which diminishes possible  
contamination with normal skin flora. All specimens  
35 were digested with Bam HI prior to hybridization.

006290\*62F20560

KS390Bam and KS480Bam hybridized nonspecifically to both KS and non-KS tissues and were not further characterized. 20 of 27 (74%) AIDS-KS DNAs hybridized with variable intensity to both KS330Bam and KS627Bam, and one additional KS specimen hybridized only to KS627Bam by Southern blotting (Figure 2 and Table 1). In contrast to AIDS-KS lesions, only 6 of 39 (15%) non-KS tissues from patients with AIDS hybridized to the KS330Bam and KS627Bam inserts (Table 1).

Specific hybridization did not occur with lymphoma or lymph node DNA from 36 persons without AIDS or with control DNA from 49 tissue biopsy specimens obtained from a consecutive series of patients. DNA extracted from several vascular tumors, including a hemangiopericytoma, two angiosarcomas and a lymphangioma, were also negative by Southern blot hybridization. DNA extracted from tissues with opportunistic infections common to AIDS patients, including 7 acid-fast bacillus (undetermined species), 1 cytomegalovirus, 1 cat-scratch bacillus, 2 cryptococcus and 1 toxoplasmosis infected tissues, were negative by Southern blot hybridization to KS330Bam and KS627Bam (Table 1).

**Table 1.** Southern blot hybridization for KS330Bam and KS627Bam and PCR amplification for KS330<sub>234</sub> in human tissues from individual patients.

<u>Tissue</u>	n	KS330Bam Southern hybridization n(%)	KS627Bam Southern hybridization n(%)	KS330 <sub>234</sub> PCR positive
AIDS-KS	27*	20 (74)	21 (78)	25 (93)
AIDS lymphomas	27†	3 (11)	3 (11)	3 (11)
AIDS lymph nodes	12	3 (25)	3 (25)	3 (25)
Non-AIDS Lymphomas	29	0 (0)	0 (0)	0 (0)
Non-AIDS lymph nodes	7	0 (0)	0 (0)	0 (0)
Vascular tumors	4§	0 (0)	0 (0)	0 (0)
Opportunistic 13Π infections		0 (0)	0 (0)	0 (0)
Consecutive surgical biopsies	49¶**	0 (0)	0 (0)	0 (0)

Legend to Table 1:

- 5 \*Includes one AIDS-KS specimen unamplifiable for p53 exon 6 and one tissue which on microscopic examination did not have any detectable KS tissue present. Both of these samples were negative by Southern blot hybridization to KS330Bam and KS627Bam and by PCR amplification for the KS330<sub>234</sub> amplicon.
- 10 †Includes 7 small non-cleaved cell lymphomas, 20 diffuse large cell and immunoblastic lymphomas. Three of the lymphomas with immunoblastic morphology were positive for KS330Bam and KS627Bam.
- 15 ‡ Includes 13 anaplastic large cell lymphomas, 4 diffuse large cell lymphomas, 4 small lymphocytic lymphomas/chronic lymphocytic leukemias, 3 hairy cell leukemias, 2 monocytoid B-cell lymphomas, 1 follicular small cleaved cell lymphoma, 1 Burkitt's lymphoma, 1  
20 plasmacytoma.
- § Includes 2 angiosarcomas, 1 hemangiopericytoma and 1 lymphangioma.
- 25 ¶ Includes 2 cryptococcus, 1 toxoplasmosis, 1 cat-scratch bacillus, 1 cytomegalovirus, 1 Epstein-Barr virus, and 7 acid-fast bacillus infected tissues. In addition, pure cultures of Mycobacterium avium-complex were negative by Southern hybridization and PCR, and  
30 pure cultures of Mycoplasma penetrans were negative by PCR.
- 35 ¶ Tissues included skin, appendix, kidney, prostate, hernia sac, lung, fibrous tissue, gallbladder, colon, foreskin, thyroid, small bowel, adenoid, vein, axillary tissue, lipoma, heart, mouth, hemorrhoid, pseudoaneurysm and fistula track. Tissues were

collected from a consecutive series of biopsies on patients without AIDS but with unknown HIV serostatus.

5     \*\*Apparent nonspecific hybridization at approximately  
20 Kb occurred in 4 consecutive surgical biopsy DNA  
samples: one colon and one hernia sac DNA sample  
hybridized to KS330Bam alone, another hernia sac DNA  
sample hybridized to KS627Bam alone and one appendix  
10    DNA sample hybridized to both KS330Bam and KS627Bam.  
These samples did not hybridize in the 330-630 bp  
range expected for these sequences and were PCR  
negative for KS330<sub>234</sub>.

15

RECEIVED OCT 24 1996

In addition, DNA from Epstein-Barr virus-infected peripheral blood lymphocytes and pure cultures of Mycobacterium avium-complex were also negative by Southern hybridization. Overall, 20 of 27 (74%) AIDS-KS specimens hybridized to KS330Bam and 21 of 27 (78%) AIDS-KS specimens hybridized to KS627Bam, compared to only 6 of 142 (4%) non-KS human DNA control specimens ( $\chi^2=85.02$ ,  $p < 10^{-7}$  and  $\chi^2=92.4$ ,  $p < 10^{-7}$  respectively).

The sequence copy number in the AIDS-KS tissues was estimated by simultaneous hybridization with KS330Bam and a 440 bp probe for the constant region of the T cell receptor  $\beta$  gene [76]. Samples in lanes 5 and 6 of Figures 2A-2B showed similar intensities for the two probes indicating an average copy number of approximately two KS330Bam sequences per cell, while remaining tissues had weaker hybridization signals for the KS330Bam probe.

### 20 Experiment 3: Characterization of KS330Bam and KS627Bam

To further characterize KS330Bam and KS627Bam, six clones for each insert were sequenced. The Sequenase version 2.0 (United States Biochemical, Cleveland, OH) system was used and sequencing was performed according to manufacturer's instructions. Nucleotide sequences were confirmed with an Applied Biosystems 373A Sequencer in the DNA Sequencing Facilities at Columbia University.

KS330Bam is a 330 bp sequence with 51% G:C content (Figure 3B) and KS627Bam is a 627 bp sequence with a 63% G:C content (Figure 3C). KS330Bam has 54% nucleotide identity to the BDLF1 open reading frame (ORF) of Epstein-Barr virus (EBV). Further analysis revealed that both KS330Bam and KS627Bam code for



amino acid sequences with homology to polypeptides of viral origin. SwissProt and PIR protein databases were searched for homologous ORF using BLASTX [3].

5 KS330Bam is 51% identical by amino acid homology to a portion of the ORF26 open reading frame encoding the capsid protein VP23 (NCBI g.i. 60348, bp 46024 - 46935) of herpesvirus saimiri [2], a gammaherpesvirus which causes fulminant lymphoma in New world monkeys.  
10 This fragment also has a 39% identical amino acid sequence to the theoretical protein encoded by the homologous open reading frame BDLF1 in EBV (NCBI g.i. 59140, bp 132403 -133307) [9]. The amino acid sequence encoded by KS627Bam is homologous with weaker  
15 identity (31%) to the tegument protein, gp140 (ORF 29, NCBI g.i. 60396, bp108782-112681) of herpesvirus saimiri.

Sequence data from KS330Bam was used to construct PCR  
20 primers to amplify a 234bp fragment designated KS330<sub>234</sub> (Figure 3B). The conditions for PCR analyses were as follows: 94°C for 2 min (1 cycle); 94°C for 1 min, 58°C for 1 min, 72°C for 1 min (35 cycles); 72°C extension for 5 min (1 cycle). Each PCR reaction used  
25 0.1 µg of genomic DNA, 50 pmoles of each primer, 1 unit of Taq polymerase, 100 µM of each deoxynucleotide triphosphate, 50 mM KCl, 10mM Tris-HCl (pH 9.0), and 0.1% Triton-X-100 in a final volume of 25 µl. Amplifications were carried out in a Perkin-  
30 Elmer 480 Thermocycler with 1-s ramp times between steps.

Although Southern blot hybridization detected the KS330Bam sequence in only 20 of 27 KS tissues, 25 of  
35 the 27 tissues were positive by PCR amplification for KS330<sub>234</sub> (Figures 4A-4B) demonstrating that KS330Bam is present in some KS lesions at levels below the

threshold for detection by Southern blot hybridization. All KS330<sub>234</sub> PCR products hybridized to a <sup>32</sup>P end-labelled 25 bp internal oligomer, confirming the specificity of the PCR (Figure 4B). Of the two AIDS-KS specimens negative for KS330<sub>234</sub>, both specimens appeared to be negative for technical reasons: one had no microscopically detectable KS tissue in the frozen sample (Figures 4A-4B, lane 3), and the other (Figures 4A-4B, lane 15) was negative in the control PCR amplification for the p53 gene indicating either DNA degradation or the presence of PCR inhibitors in the sample. PCR amplification of the p53 tumor suppressor gene was used as a control for DNA quality. Sequences of p53 primers from P6-5, 5'-ACAGGGCTGGTTGCCAGGGT-3' (SEQ ID No: 44); and P6-3. 5'-AGTTGCAAACCAGACCTCAG-3' (SEQ ID NO: 45) [25].

Except for the 6 control samples from AIDS patients that were also positive by Southern blot hybridization, none of the other 136 control specimens were positive by PCR for KS330<sub>234</sub>. All of these specimens were amplifiable for the p53 gene, indicating that inadequate PCR amplification was not the reason for lack of detection of KS330<sub>234</sub> in the control tissues. Samples containing DNA from two candidate KS agents, EBV and Mycoplasma penetrans (ATCC Accession No. 55252), a pathogen commonly found in the genital tract of patients with AIDS-KS [59] were also negative for amplification of KS330<sub>234</sub>. In addition, several KS specimens were tested using commercial PCR primers (Stratagene, La Jolla, CA) specific for mycoplasmata and primers specific for the EBNA-2, EBNA-3C and EBER regions of EBV and were negative [57].

Overall, DNA from 25 (93%) of 27 AIDS-KS tissues were positive by PCR compared with DNA from 6 (4%) of 142

control tissues, including 6 (15%) of 39 non-KS lymph nodes and lymphomas from AIDS patients ( $\chi^2=38.2$ ,  $p < 10^{-6}$ ), 0 of 36 lymph nodes and lymphomas from nonAIDS patients ( $\chi^2=55.2$ ,  $p < 10^{-7}$ ) and 0 of 49 consecutive biopsy specimens ( $\chi^2=67.7$ ,  $p < 10^{-7}$ ). Thus, KS330<sub>234</sub> was found in all 25 amplifiable tissues with microscopically detectable AIDS-KS, but rarely occurred in non-KS tissues, including tissues from AIDS patients.

10

Of the six control tissues from AIDS patients that were positive by both PCR and Southern hybridization, two patients had KS elsewhere, two did not develop KS and complete clinical histories for the remaining two patients were unobtainable. Three of the six positive non-KS tissues were lymph nodes with follicular hyperplasia taken from patients with AIDS. Given the high prevalence of KS among patients with AIDS, it is possible that undetected microscopic foci of KS were present in these lymph nodes. The other three positive tissue specimens were B cell immunoblastic lymphomas from AIDS patients. It is possible that the putative KS agent is also a cofactor for a subset of AIDS-associated lymphomas [16, 17, 80].

25

To determine whether KS330Bam and KS627Bam are portions of a larger genome and to determine the proximity of the two sequences to each other, samples of KS DNA were digested with Pvu II restriction enzymes. Digested genomic DNA from three AIDS-KS samples were hybridized to KS330Bam and KS627Bam by Southern blotting (Figure 5). These sequences hybridized to various sized fragments of the digested KS DNA indicating that both sequences are fragments of larger genomes. Differences in the KS330Bam hybridization pattern to Pvu II digests of the three AIDS-KS specimens indicate that polymorphisms may

35

006250\*62720950

occur in the larger genome. Individual fragments from the digests failed to simultaneously hybridize with both KS330Bam and KS627Bam, demonstrating that these two Bam HI restriction fragments are not adjacent to one another.

If KS330Bam and KS627Bam are heritable polymorphic DNA markers for KS, these sequences should be uniformly detected at non-KS tissue sites in patients with AIDS-KS. Alternatively, if KS330Bam and KS627Bam are sequences specific for an exogenous infectious agent, it is likely that some tissues are uninfected and lack detectable KS330Bam and KS627Bam sequences. DNA extracted from multiple uninvolved tissues from three patients with AIDS-KS were hybridized to <sup>32</sup>P-labelled KS330Bam and KS627Bam probes as well as analyzed by PCR using the KS330<sub>234</sub> primers (Table 2). While KS lesion DNA samples were positive for both bands, uninvolved tissues were frequently negative for these sequences. KS lesions from patients A, B and C, and uninvolved skin and muscle from patient A were positive for KS330Bam and KS627Bam, but muscle and brain tissue from patient B and muscle, brain, colon, heart and hilar lymph node tissues from patient C were negative for these sequences. Uninvolved stomach lining adjacent to the KS lesion in patient C was positive by PCR, but negative by Southern blotting which suggests the presence of the sequences in this tissue at levels below the detection threshold for Southern blotting.

**Table 2:** Differential detection of KS330Bam, KS627Bam and KS330<sub>234</sub> sequences in KS-involved and non-involved tissues from three patients with AIDS-KS.

	KS330Bam	KS627Bam	KS330 <sub>234</sub>
Patient A			
KS, skin	+	+	+
nl skin	+	+	+
nl muscle	+	+	+
Patient B			
KS, skin	+	+	+
nl muscle	-	-	-
nl brain	-	-	-
Patient C			
KS, stomach	+	+	+
nl stomach adjacent to KS	-	-	+
nl muscle	-	-	-
nl brain	-	-	-
nl colon	-	-	-
nl heart	-	-	-
nl hilar lymph nodes	-	-	-

**Experiment 4: Subcloning and sequencing of KSHV**

5 KS330Bam and KS627Bam are genomic fragments of a novel infectious agent associated with AIDS-KS. A genomic library from a KS lesion was made and a phage clone with a 20 kb insert containing the KS330Bam sequence was identified. The 20 kb clone digested with PvuII (which cuts in the middle of the KS330Bam sequence) produced 1.1 kb and 3 kb fragments that hybridized to  
 10 KS330Bam. The 1.1 kb subcloned insert and ~900 bp from the 3 kb subcloned insert resulting in 9404 bp of

contiguous sequence was entirely sequenced. This sequence contains partial and complete open reading frames homologous to regions in gamma herpesviruses.

5 The KS330Bam sequence is an internal portion of an 918  
bp ORF with 55-56% nucleotide identity to the ORF26  
and BDLF1 genes of HSVSA and EBV respectively. The  
EBV and HSVSA translated amino acid sequences for  
these ORFs demonstrate extensive homology with the  
10 amino acid sequence encoded by the KS-associated 918  
bp ORF (Figure 6). In HSVSA, the VP23 protein is a  
late structural protein involved in capsid  
construction. Reverse transcriptase (RT)-PCR of mRNA  
from a KS lesion is positive for transcribed KS330Bam  
15 mRNA and that indicates that this ORF is transcribed  
in KS lesions. Additional evidence for homology  
between the KS agent and herpesviruses comes from a  
comparison of the genomic organization of other  
potential ORFs on the 9404 bp sequence (Figure 3A)  
20 The 5' terminus of the sequence is composed  
nucleotides having 66-67% nucleotide identity and 68-  
71% amino acid identity to corresponding regions of  
the major capsid protein (MCP) ORFs for both EBV and  
HSVSA. This putative MCP ORF of the KS agent lies  
25 immediately 5' to the BDLF1/ORF26 homolog which is a  
conserved orientation among herpesvirus subfamilies  
for these two genes. At the 3' end of this sequence,  
the reading frame has strong amino acid and nucleotide  
homology to HSVSA ORF 27. Thus, KS-associated DNA  
30 sequences at four loci in two separate regions with  
homologies to gamma herpesviral genomes have been  
identified.

35 In addition to fragments obtained from Pvu II digest  
of the 21 Kb phage insert described above, fragments  
obtained from a BamHI/NotI digest were also subcloned  
into pBluescript (Stratagene, La Jolla, CA). The

SUB B4 7

termini of these subcloned fragments were sequenced and were also found to be homologous to nucleic acid sequence EBV and HSVSA genes. These homologs have been used to develop a preliminary map of subcloned fragments (Figure 9). Thus, sequencing has revealed that the KS agent maintains co-linear homology to gamma herpesviruses over the length of the 21 Kb phage insert.

**10     Experiment 5: Determination of the phylogeny of KSHV**

Regions flanking KS330Bam were sequenced and characterized by directional walking. This was performed by the following strategy: 1) KS genomic libraries were made and screened using the KS330Bam fragment as a hybridization probe, 2) DNA inserts from phage clones positive for the KS330Bam probe were isolated and digested with suitable restriction enzyme(s), 3) the digested fragments were subcloned into pBluescript (Stratagene, La Jolla, CA), and 4) the subclones were sequenced. Using this strategy, the major capsid protein (MCP) ORF homolog was the first important gene locus identified. Using sequenced unique 3' and 5' end-fragments from positive phage clones as probes, and following the strategy above a KS genomic library are screened by standard methods for additional contiguous sequences.

For sequencing purposes, restriction fragments are subcloned into phagemid pBluescript KS+, pBluescript KS-, pBS+, or pBS- (Stratagene) or into plasmid pUC18 or pUC19. Recombinant DNA was purified through CsCl density gradients or by anion-exchange chromatography (Qiagen).

35

Nucleotide sequenced by standard screening methods of cloned fragments of KSHV were done by direct

sequencing of double-stranded DNA using oligonucleotide primers synthesized commercially to "walk" along the fragments by the dideoxy-nucleotide chain termination method. Junctions between clones  
5 are confirmed by sequencing overlapping clones.

Targeted homologous genes in regions flanking KS330Bam include, but are not limited to: Il-10 homolog, thymidine kinase (TK), g85, g35, gH, capsid proteins and MCP. TK is an early protein of the herpesviruses  
10 functionally linked to DNA replication and a target enzyme for anti-herpesviral nucleosides. TK phosphorylates acyclic nucleosides such as acyclovir which in turn inhibit viral DNA polymerase chain  
15 extension. Determining the sequence of this gene will aid in the prediction of chemotherapeutic agents useful against KSHV. TK is encoded by the EBV BRLF1 ORF located -9700 bp rightward of BDLF1 and by the HSVSA ORF 21 -9200 bp rightward of the ORF 26. A  
20 subcloned fragment of KS5 was identified with strong homology to the EBV and HSVSA TK open reading frames.

g85 is a late glycoprotein involved in membrane fusion homologous to gH in HSV1. In EBV, this protein is  
25 encoded by BLXF2 ORF located -7600 bp rightward of BDLF1, and in HSVSA it is encoded by ORF 22 located -7100 bp rightward of ORF26.

g35 is a late EBV glycoprotein found in virion and plasma membrane. It is encoded by BDLF3 ORF which is  
30 1300 bp leftward of BDLF1 in EBV. There is no BDLF3 homolog in HSVSA. A subcloned fragment has already been identified with strong homology to the EBV gp35 open reading frame.

35 Major capsid protein (MCP) is a conserved 150 KDa protein which is the major component of herpesvirus

006250 6474960



capsid. Antibodies are generated against the MCP during natural infection with most herpesviruses. The terminal 1026 bp of this major capsid gene homolog in KSHV have been sequenced.

5

Targeted homologous genes/loci in regions flanking KS627Bam include, but are not limited to: terminal reiterated repeats, LMPI, EBERs and Ori P. Terminal reiterated sequences are present in all herpesviruses. In EBV, tandemly reiterated 0.5 Kb long terminal repeats flank the ends of the linear genome and become joined in the circular form. The terminal repeat region is immediately adjacent to BNRF1 in EBV and ORF 75 in HSVSA. Since the number of terminal repeats varies between viral strains, identification of terminal repeat regions may allow typing and clonality studies of KSHV in KS lesions. Sequencing through the terminal repeat region may determine whether this virus is integrated into human genome in KS.

10

LMPI is an latent protein important in the transforming effects of EBV in Burkitt's lymphoma. This gene is encoded by the EBV BNRF1 ORF located ~2000 bp rightward of tegument protein ORF BNRF1 in the circularized genome. There is no LMP1 homolog in HSVSA.

15

EBERs are the most abundant RNA in latently EBV infected cells and Ori-P is the origin of replication for latent EBV genome. This region is located between -4000-9000 bp leftward of the BNRF1 ORF in EBV; there are no corresponding regions in HSVSA.

20

The data indicates that the KS agent is a new human herpesvirus related to gamma herpesviruses EBV and HSVSA. The results are not due to contamination or to incidental co-infection with a known herpesvirus since

25

the sequences are distinct from all sequenced herpesviral genomes (including EBV, CMV, HHV6 and HSVSA) and are associated specifically with KS in three separate comparative studies. Furthermore, PCR testing of KS DNA with primers specific for EBV-1 and EBV-2 failed to demonstrate these viral genomes in these tissues. Although KSHV is homologous to EBV regions, the sequence does not match any other known sequence and thus provides evidence for a new viral genome, related to but distinct from known members of the herpesvirus family.

**Experiment 6: Serological studies**

**Indirect immunofluorescence assay (IFA)**

Virus-containing cells are coated to a microscope slide. The slides are treated with organic fixatives, dried and then incubated with patient sera. Antibodies in the sera bind to the cells, and then excess nonspecific antibodies are washed off. An antihuman immunoglobulin linked to a fluorochrome, such as fluorescein, is then incubated with the slides, and then excess fluorescent immunoglobulin is washed off. The slides are then examined under a microscope and if the cells fluoresce, then this indicates that the sera contains antibodies directed against the antigens present in the cells, such as the virus.

An indirect immunofluorescence assay (IFA) was performed on the Body Cavity-Based Lymphoma cell line (BCBL-1), which is a naturally transformed EBV infected (nonproducing) B cell line, using 4 KS patient sera and 4 control sera (from AIDS patients without KS). Initially, both sets of sera showed similar levels of antibody binding. To remove

nonspecific antibodies directed against EBV and lymphocyte antigens, sera at 1:25 dilution were pre-adsorbed using  $3 \times 10^6$  1% paraformaldehyde-fixed Raji cells per ml of sera. BCBL1 cells were fixed with ethanol/acetone, incubated with dilutions of patient sera, washed and incubated with fluorescein-conjugated goat anti-human IgG. Indirect immunofluorescent staining was determined.

Table 3 shows that unadsorbed case and control sera have similar end-point dilution indirect immunofluorescence assay (IFA) titers against the BCBL1 cell line. After Raji adsorption, case sera have four-fold higher IFA titers against BCBL1 cells than control sera. Results indicated that pre-adsorption against paraformaldehyde-fixed Raji cells reduces fluorescent antibody binding in control sera but do not eliminate antibody binding to KS case sera. These results indicate that subjects with KS have specific antibodies directed against the KS agent that can be detected in serological assays such as IFA, Western blot and Enzyme immunoassays (Table 3).

006290"64F20960

**Table 3: Indirect immunofluorescence end-point titers for KS case and non-KS control sera against the BCBL-1 cell line**

	<u>Sera No.</u>	<u>Status*</u>	<u>Pre-adsorption</u>	<u>Post-adsorption**</u>
5	1	KS	≥ 1:400	≥ 1:400
	2	KS	1:100	1:100
	3	KS	1:200	1:100
10	4	KS	≥ 1:400	1:200
	5	Control	≥ 1:400	1:50
	6	Control	1:50	1:50
	7	Control	1:100	1:50
15	8	Control	1:200	1:50

**Legend Table 3:**

- 20 \* KS=autopsy-confirmed male, AIDS patient  
Control=autopsy-confirmed female, AIDS patient, no KS
- 25 \*\* Adsorbed against RAJI cells treated with 1% paraformaldehyde

Immunoblotting ("Western blot")

30 Virus-containing cells or purified virus (or a portion of the virus, such as a fusion protein) is electrophoresed on a polyacrylamide gel to separate the protein antigens by molecular weight. The proteins are blotted onto a nitrocellulose or nylon membrane, then the membrane is incubated in patient

35 sera. Antibodies directed against specific antigens are developed by incubating with a anti-human immunoglobulin attached to a reporter enzyme, such as a peroxidase. After developing the membrane, each antigen reacting against antibodies in patient sera

40 shows up as a band on the membrane at the corresponding molecular weight region.

Enzyme immunoassay ("EIA or ELISA")

5 Virus-containing cells or purified virus (or a portion  
of the virus, such as a fusion protein) is coated to  
the bottom of a 96-well plate by various means  
(generally incubating in alkaline carbonate buffer).  
The plates are washed, then the wells are incubated  
with patient sera. Antibodies in the sera directed  
against specific antigens stick on the plate. The  
10 wells are washed again to remove nonspecific antibody,  
then they are incubated with a antihuman  
immunoglobulin attached to a reporter enzyme, such as  
a peroxidase. The plate is washed again to remove  
nonspecific antibody and then developed. Wells  
15 containing antigen that is specifically recognized by  
antibodies in the patients sera change color and can  
be detected by an ELISA plate reader (a  
spectrophotomer).

20 All three of these methods can be made more specific  
by pre-incubating patient sera with uninfected cells  
to adsorb out cross-reacting antibodies against the  
cells or against other viruses that may be present in  
the cell line, such as EBV. Cross-reacting antibodies  
25 can potentially give a falsely positive test result  
(i.e. the patient is actually not infected with the  
virus but has a positive test result because of cross-  
reacting antibodies directed against cell antigens in  
the preparation). The importance of the infection  
30 experiments with Raji is that if Raji cells, or  
another well-defined cell line, can be infected, then  
the patient's sera can be pre-adsorbed against the  
uninfected parental cell line and then tested in one  
of the assays. The only antibodies left in the sera  
35 after pre-adsorption that bind to antigens in the  
preparation should be directed against the virus.

Experiment 7:

5 BCBL 1, from lymphomatous tissues belonging to a rare  
infiltrating, anaplastic body cavity lymphoma  
occurring in AIDS patients has been placed in  
continuous cell culture and shown to be continuously  
infected with the KS agent. This cell line is also  
naturally infected with Epstein-Barr Virus (EBV). The  
10 BCBL cell line was used as an antigen substrate to  
detect specific KS antibodies in persons infected with  
the putative virus by Western-blotting. Three  
lymphoid B cell lines were used as controls. These  
included the EBV genome positive cell line P3H3, the  
15 EBV genome defective cell line Raji and the EBV genome  
negative cell line Bjab.

Cells from late-log phase culture were washed 3 time  
with PBS by centrifugation at 500 g for 10min. and  
20 suspended in sample buffer containing 50 mM Tris-HCl  
pH 6.8, 2% SDS (w/v), 15% glycerol (v/v), 5%  $\beta$ -  
mercaptoethanol (v/v) and 0.001% bromophenol (w/v)  
with protease inhibitor, 100  $\mu$ M phenylmethylsulfonyl  
fluoride (PMSF). The sample was boiled at 100°C for  
25 5 min and centrifuged at 14,000 g for 10 min. The  
proteins in the supernatant was then fractionated by  
sodium, dodecyl sulfate-polyacrylamide gel  
electrophoresis (SDS-PAGE) under reducing conditions  
with a separation gel of 15% and a stacking gel of 5%  
30 (3). Prestained protein standards were included:  
myosin, 200 kDa;  $\beta$ -galactosidase, 118 kDa; BSA, 78  
kDa; ovalbumin, 47.1 kDa; carbonic anhydrase, 31.4  
kDa; soybean trypsin inhibitor, 25.5 kDa, lysozyme,  
18.8 kDa and aprotinin, 8.3 kDa (Bio-Rad).  
35 Immunoblotting experiments were performed according to  
the method of Towbin et al. (4). Briefly, the  
proteins were electrophoretically transferred to

Hybon-C extra membranes (Pharmacia) at 24 V for 70 min. The membranes were then dried at 37°C for 30 min, saturated with 5% skim milk in Tris-buffered saline, pH 7.4 (TBS) containing 50 mM Tris-HCl and 200 mM NaCl, at room temperature for 1 h. The membranes were subsequently incubated with human sera at dilution 1:200 in 1% skim milk overnight at room temperature, washed 3 times with a solution containing TBS, 0.2% Triton X-100 and 0.05% skim milk and then 2 times with TBS. The membranes were then incubated for 2 h at room temperature with alkaline phosphatase conjugated goat anti-mouse IgG + IgM + IgA (Sigma) diluted at 1:5000 in 1% skim milk. After repeating the washing, the membranes were stained with nitroblue tetranolium chloride and 5-bromo-4-chloro-3-indolylphosphate p-toluidine salt (Gibco BRL).

Two bands of approximately 226 kDa and 234 kDa were identified to be specifically present on the Western blot of BCBL cell lysate in 5 sera from AIDS gay man patients infected with KS. These 2 bands were absent from the lysates of P3H3, Raji and Bjab cell lysates. 5 sera from AIDS gay man patients without KS and 2 sera from AIDS woman patients without KS as well as 1 sera from nasopharyngeal carcinoma patient were not able to detect these 2 bands in BCBL 1, P3H3, Raji and Bjab cell lysates. In a blinded experiment, using the 226 kDa and 234 kDa markers, 15 out of 16 sera from KS patients were correctly identified. In total, the 226 kDa and 234 kDa markers were detected in 20 out of 21 sera from KS patients.

The antigen is enriched in the nuclei fraction of BCBL1. Enriched antigen with low background can be obtained by preparing nucleic from BCBC as the starting antigen preparation using standard, widely available protocols. For example, 500-750ml of BCBL

at  $5 \times 10^5$  cells/ml can be pelleted at low speed. The pellet is placed in 10 mM NaCl, 10 mM Tris pH 7.8, 1.5 mM MgCl<sub>2</sub> (equi volume) + 1.0% NP-40 on ice for 20 min to lyse cells. The lysate is then spun at 1500 rpm for 10 min. to pellet nucleic. The pellet is used as the starting fraction for the antigen preparation for the Western blot. This will reduce cross-reactive cytoplasmic antigens.

10 **Experiment 8: Transmission studies**

**Co-infection experiments**

BCBL1 cells were co-cultivated with Raji cell lines separated by a 0.45  $\mu$  tissue filter insert. Approximately,  $1-2 \times 10^6$  BCBL1 and  $2 \times 10^6$  Raji cells were co-cultivated for 2-20 days in supplemented RPMI alone, in 10  $\mu$ g/ml 5'-bromodeoxyuridine (BUdR) and 0.6  $\mu$ g/ml 5'-flourodeoxyuridine or 20 ng/ml 12-O-tetradecanoylphorbol-13-acetate (TPA). After 2, 8, 12 or 20 days co-cultivation, Raji cells were removed, washed and placed in supplemented RPMI 1640 media. A Raji culture co-cultivated with BCBL1 in 20 ng/ml TPA for 2 days survived and has been kept in continuous suspension culture for >10 weeks. This cell line, designated RCC1 (Raji Co-Culture, No. 1) remains PCR positive for the KS330<sub>234</sub> sequence after multiple passages. This cell line is identical to its parental Raji cell line by flow cytometry using EMA, B1, B4 and BerH2 lymphocyte-flow cytometry (approximately 2%). RCC1 periodically undergo rapid cytolysis suggestive of lytic reproduction of the agent. Thus, RCC1 is a Raji cell line newly infected with KSHV.

35 The results indicate the presence of a new human virus, specifically a herpesvirus in KS lesions. The high degree of association between this agent and



AIDS-KS (>90%), and the low prevalence of the agent in non-KS tissues from immunocompromised AIDS patients, indicates that this agent has a causal role in AIDS-KS [47, 68].

5

Experiment 10<sup>9</sup>: Isolation of KSHV

Crude virus preparations are made from either the supernatant or low speed pelleted cell fraction of BCBL1 cultures. Approximately 650ml or more of log phase cells should be used (>5X10<sup>6</sup> cells/ml).

For bonding whole virion from supernatant, the cell free supernatant is spun at 10,000 rpm in a GSA rotor for 10 min to remove debris. PEG-8000 is added to 7%, dissolved and placed on ice for >2.5 hours. The PEG-supernatant is then spun at 10,000 xg for 30 min. supernatant is poured off and the pellet is dried and scraped together from the centrifuge bottles. The pellet is then resuspended in a small volume (1-2 ml) of virus buffer (VB, 0.1 M NaCl, 0.01 M Tris, pH 7.5). This procedure will precipitate both naked genome and whole virion. The virion are then isolated by centrifugation at 25,000 rpm in a 10-50% sucrose gradient made with VB. One ml fractions of the gradient are then obtained by standard techniques (e.g. using a fractionator) and each fraction is then tested by dot blotting using specific hybridizing primer sequences to determine the gradient fraction containing the purified virus (preparation of the fraction maybe needed in order to detect the presence of the virus, such as standard DNA extraction).

To obtain the episomal DNA from the virus, the pellet of cells is washed and pelleted in PBS, then lysed using hypotonic shock and/or repeated cycles of freezing and thawing in a small volume (<3 ml).

Nuclei and other cytoplasmic debris are removed by centrifugation at 10,000g for 10 min, filtration through a 0.45 m filter and then repeat centrifugation at 10,000g for 10 min. This crude preparation contains viral genome and soluble cell components. The genome preparation can then be gently chloroform-phenol extracted to remove associated proteins or can be placed in neutral DNA buffer (1 M NaCl, 50 mM Tris, 10 mM EDTA, pH 7.2-7.6) with 2% sodium dodecylsulfate (SDS) and 1% sarcosyl. The genome is then banded by centrifugation through 10-30% sucrose gradient in neutral DNA buffer containing 0.15% sarcosyl at 20,000 rpm in a SW 27.1 rotor for 12 hours (for 40,000 rpm for 2-3 hours in an SW41 rotor). The band is detected as described above.

An example of the method for isolating KSHV genome from KSHV infected cell cultures (97 and 98). Approximately 800 ml of BCBL1 cells are pelleted, washed with saline, and pelleted by low speed centrifugation. The cell pellet is lysed with an equal volume of RSB (10 mM NaCl, 10 mM Tris-HCl, 1.5 mM MgCl<sub>2</sub>, pH 7.8) with 1% NP-40 on ice for 10 minutes. The lysate is centrifuged at 900xg for 10 minutes to pellet nuclei. This step is repeated. To the supernatant is added 0.4% sodium dodecylsulfate and EDTA to a final concentration of 10 mM. The supernatant is loaded on a 10-30% sucrose gradient in 1.0 M NaCl, 1mM EDTA, 50mM Tris-HCl, pH 7.5. The gradients are centrifuged at 20,000 rpm on a SW 27.1 rotor for 12 hours. In figure 11, 0.5 ml aliquots of the gradient have been fractionated (fractions 1-62) with the 30% gradient fraction being at fraction No. 1 and the 10% gradient fraction being at fraction No. 62. Each fraction has been dot hybridized to a nitrocellulose membrane and then a <sup>32</sup>P-labeled KSHV DNA fragment, KS631Bam has been hybridized to the membrane

using standard techniques. Figure 11 shows that the major solubilized fraction of the KSHV genome bands (i.e. is isolated) in fractions 42 through 48 of the gradient with a high concentration of the genome being present in fraction 44. A second band of solubilized KSHV DNA occurs in fractions 26 through 32.

Experiment 10: Purification of KSHV

DNA is extracted using standard techniques from the RCC-1 or RCC-1<sub>2FS</sub> cell line [27, 49, 66]. The DNA is tested for the presence of the KSHV by Southern blotting and PCR using the specific probes as described hereinafter. Fresh lymphoma tissue containing viable infected cells is simultaneously filtered to form a single cell suspension by standard techniques [49, 66]. The cells are separated by standard Ficoll-Plaque centrifugation and lymphocyte layer is removed. The lymphocytes are then placed at  $>1 \times 10^6$  cells/ml into standard lymphocyte tissue culture medium, such as RMP 1640 supplemented with 10% fetal calf serum. Immortalized lymphocytes containing the KSHV virus are indefinitely grown in the culture media while nonimmortalized cells die during course of prolonged cultivation.

Further, the virus may be propagated in a new cell line by removing media supernatant containing the virus from a continuously infected cell line at a concentration of  $>1 \times 10^6$  cells/ml. The media is centrifuged at 2000xg for 10 minutes and filtered through a  $0.45 \mu$  filter to remove cells. The media is applied in a 1:1 volume with cells growing at  $>1 \times 10^6$  cells/ml for 48 hours. The cells are washed and pelleted and placed in fresh culture medium, and tested after 14 days of growth.

The herpesvirus may be isolated from the cell DNA in the following manner. An infected cell line, which can be lysed using standard methods such as hyposmotic shocking and Dounce homogenization, is first pelleted at 2000xg for 10 minutes, the supernatant is removed and centrifuged again at 10,000xg for 15 minutes to remove nuclei and organelles. The supernatant is filtered through a 0.45 $\mu$  filter and centrifuged again at 100,000xg for 1 hour to pellet the virus. The virus can then be washed and centrifuged again at 100,000xg for 1 hour.

005230\*6474956

REFERENCES:

1. Ablashi, D.V., et al. Virology 184:545-552.
2. Albrecht, J.C., et al. (1992) J. Virol. 66:5047.
- 5 3. Altshul, S.F., et al. (1990) J. Molec. Biol. 215:403.
4. Analytical Biochemistry (1984) 238:267-284.
5. Andrei, et al. (1992) Eur. J. Clin. Microbiol. Infect. Dis. 11(2):143-51.
- 10 6. Archibald, C.P., et al. (1992) Epidemiol. 3:203.
7. Asada, H., et al (1989) J. Clin. Microbiol. 27(10):2204.
8. Ausubel, F., et al. (1987) Current Protocols in Molecular Biology, New York.
- 15 9. Baer, R.J., et al. (1984) Nature 310:207.
10. Bagasra, et al. (1992) J. New England Journal of Medicine 326(21):1385-1391.
11. Balzarini, et al. (1990) Mol. Pharm. 37,402-7.
12. Basic and Clinical Immunology 7th Edition D. Stites and A. Terr ed.
- 20 13. Beral, V., et al. (1990) Lancet 335:123.
14. Beral, V., et al. (1991) Brit. Med. J. 302:624.
15. Beral, V., et al. (1992) Lancet 339:632.
16. Bendsøe, N., et al. (1990) Eur. J. Cancer 26:699.
- 25 17. Biggar, R.J., et al. (1994) Am. J. Epidemiol. 139:362.
18. Bovenzi, P., et al. (1993) Lancet 341:1288.
19. Beaucage and Carruthers (1981) Tetrahedron Lett. 22:1859-1862.
- 30 20. Braitman, et al. (1991) Antimicrob. Agents and Chemotherapy 35(7):1464-8.
21. Burns and Sanford, (1990) J. Infect. Dis. 162(3):634-7.
22. De Clercq, (1993) Antimicrobial Chemotherapy 32, Suppl. A, 121-132.
- 35 23. Drew, W.L., et al. (1982) Lancet ii:125.
24. Falk, et al. (1991) Nature 351:290.

25. Gaidano, G., et al. (1991) Proc. Natl. Acad. Sci. USA 88:5413.
26. Gershon, A.A., (1992) J. Inf. Des. 166(Suppl):563.
- 5 27. Glick, J.L., (1980) Fundamentals of Human Lymphoid Culture, Marcel Dekker, New York.
28. Gorbach, S.L., et al. (1992) Infectious Disease Ch.35:289, W.B. Saunders, Philadelphia, Pennsylvania.
- 10 29. Greenspan, et al. (1990) J. Acquir. Immune Defic. Syndr. 3 (6):571.
30. Hardy, I., et al. (1990) Inf. Dis. Clin. N. Amer. 4(1):159.
31. Hardy, I., et al. (1991) New Engl. J. Med. 325 (22):1545.
- 15 31A. Harel-Bellan, A., et al. (1988) Exp. Med. 168:2309-2318
32. Harlow and Lane, (1988) Antibodies, A Laboratory Manual, Cold Spring Harbor Publication, New York.
- 20 33. Haverkos, H.W., et al. (1985) Sexually Transm. Dis. 12:203.
34. Helene, C. and Toulme, J. (1990) Biochim. Biophys. Acta. 1049:99-125.
35. Heniford, et al. (1993) Nucleic Acids Research 21(14):3159-3166.
- 25 36. Higashi, K., et al. (1989) J. Clin. Micro. 27(10):2204.
37. Holmberg, S.D., et al. (1990) Cancer Detection and Prevention 14:331.
- 30 38. Holliday, J., and Williams, M.V., (1992) Antimicrob. Agents Chemother. 36(9):1935.
39. Hoogenboom, H.R., et al. (1991) Nuc. Acids Res. 19:4133.
40. Hunt, et al. (1991) Eur. J. Immunol. 21:2963-2970.
- 35 41. Hybridization of Nucleic Acids Immobilized on Solid Supports Meinkoth, J. and Wahl, G.

1996-04-25

42. Hybridization with Nucleic Acid Probes pp. 495-524, (1993) Elsevier, Amsterdam.
43. Ickes, et al. (1994) Antiviral Research 23, Seventh International Conf. on Antiviral Research, Abstract No. 122, Supp. 1.
- 5 44. Jahan, N., et al. (1989) AIDS Research and Human Retroviruses 5:225.
45. Jardetzkey, et al. (1991) Nature 353:326.
46. Johnston, G.S., et al. (1990) Cancer Detection and Prevention 14:337.
- 10 47. Jung, J.U., et al. (1991) Proc. Natl. Acad. Sci. USA 88:7051.
48. Kikuta, et al. (1989) Lancet Oct. 7:861.
49. Knowles, D.M., et al. (1989) Blood 73:792-798.
- 15 50. Kohler and Milstein, (1976) Eur. J. Immunol. 6:511-519.
51. Kucera, et al. (1993) AIDS Res. Human Retroviruses 9:307-314.
52. Laboratory Techniques in Biochemistry and Molecular Biology (1978) North Holland Publishing Company, New York.
- 20 53. Lasky, L.A., (1990) J. Med. Virol. 31(1):59.
54. Levin, M.J., et al. (1992) J. Inf. Dis. 166(2):253.
- 25 55. Lifson, A.R., et al. (1990) Am. J. Epidemiol. 131:221.
56. Lin, et al. (1991) Antimicrob Agents Chemother 35(11):2440-3.
57. Lin, J.C., et al. (1993) Blood 81:3372.
- 30 58. Lisitsyn, N., et al. (1993) Science 259:946.
59. Lo, S -C., et al. (1992) Internat. J. Systematic Bacteriol. 42:357.
60. Marks, J.D., et al. (1991) J. Mol Biol. 222:581-597.
- 35 61. Marloes, et al. (1991) Eur. J. Immunol. 21:2963-2970.
62. Matteucci, et al. (1981) Am. Chem. Soc. 103:3185.

00000000000000000000

63. Maxam, A.M. and Gilbert, W. Methods in Enzymology (1980) Grossman, L. and Moldave, D., eds., Academic Press, New York, 65:499-560.
64. McCafferty, J., et al. (1990) Nature 348:552.
- 5 65. Means and Feeney, (1990) Bioconjugate Chem. A recent review of protein modification techniques, 1:2-12.
66. Metcalf, D. (1984) Clonal Culture of Hematopoietic Cells: Techniques and Applications, Elvier, New York.
- 10 67. Methods in Enzymology Vol. 152, (1987) Berger, S. and Kimmel, A. ed., Academic Press, New York
68. Miller, G., Virology (1990) B. N. Fields, D.M. Knipe eds., Raven Press, New York, 2:1921.
- 15 69. Needham-VanDevanter, D.R., et al., (1984) Nucleic Acids Res. 12:6159-6168.
70. Needleman and Wunsch, (1970) J. Mol. Biol. 48:443.
71. Neuvo, et al. (1993) American Journal of Surgical Pathology 17(7), 683-690.
- 20 72. Nucleic Acid Hybridization: A Practical Approach (1985) Ed. Hames, B.D. and Higgins, S.J., IRL Press.
73. Oren and Soble, (1991) Clinical Infectious Diseases 14:741-6.
- 25 74. PCR Protocols: A Guide to Methods and Applications. (1990) Innis, M., Gelfand D., Sninsky, J. and White, T., eds., Academic Press, San Diego.
- 30 75. Pearson and Lipman, (1988) Proc. Natl. Acad. Sci. (USA) 85:2444.
- 75A. Pearson, J.D., and Regnier, F.E., (1983) J. Chrom. 255:137-14976.
76. Pellici, P.G., et al. (1985) J. Exp. Med. 162:1015.
- 35 77. Peterman, T.A., et al. (1991) Cancer Surveys Imperial Cancer Research Fund, London, 10:23-37.

ORIGINAL PAGE



78. Roizman, B. (1991) Rev. Inf. Disease 13 Suppl. 11:S892.
79. Röttschke and Falk, (1991) Immunol. Today 12:447.
80. Safai, B., et al. (1980) Cancer 45:1472.
- 5 81. Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2nd ed.), Cold Spring Harbor Laboratory, Vols. 1-3.
82. Saunders, et al. (1990) J. Acquir. Immune Defic. Syndr. 3 (6):571.
- 10 83. Schecter, M.T., et al. (1991) Am. J. Epidemiol. 134:485.
84. Scopes, R., (1982) Protein Purification: Principles and Practice Springer-Verlag, New York.
- 15 85. Siddiqui, A., et al. (1983) Proc. Natl. Acad. Sci. USA 80:4861.
86. Skinner, G.R., et al. (1991) Comp. Immuno. Microbiol. Inf. Dis. 14(2):13.
87. Skinner, G.R., et al. (1992) Med. Microbiol. Immunol. 180(6):305.
- 20 Smith and Waterman (1981) Adv. Appl. Math. 2:482.
88. Snoeck, et al. (1992) Eur. J. Clin. Micro. Infect. Dis. 11(12):1144-55.
89. Stals, et al. (1993) Antimicrobial Agents Chemother. 37(2):218-23.
- 25 90. van den Berg, F. et al. (1989) J. Clin. Pathol. 42:128.
91. Vogel, J., et al. (1988) Nature 335:606.
92. Wang, R.H. -Y., et al. (1993) Clin. Infect. Dis. 17:724.
- 30 93. Wickstrom, E.L., et al. (1988) PNAS (USA) 85:1028-1032.
94. Winkelmann, et al. (1988) Drug Res. 38, 1545-48.
95. Winkler, et al. (1990) Antiviral Research 14:61-74.
- 35 96. Yamandaka, et al. (1991) Mol. Pharmacol. 40(3):446.

H06250-627096

97. Pellicer, A. et al. (1978) Cell 14:133-141.
98. Gibson, W. and Roizmann B. (1972) J. Virol.  
10:1044-52.

005293 6272968

**EXPERIMENTAL DETAILS SECTION II:**

Sequencing Studies: A lambda phage (KS5) from a KS  
lesion genomic library identified by positive  
5 hybridization with KS330Bam was digested with BamHI  
and Not I (Boehringer-Mannheim, Indianapolis IN); five  
fragments were gel isolated and subcloned into  
Bluescript II KS (Stratagene, La Jolla CA). The  
entire sequence was determined by bidirectional  
10 sequencing at a seven fold average redundancy by  
primer walking and nested deletions.

DNA sequence data were compiled and aligned using  
ALIGN (IBI-Kodak, Rochester NY) and analyzed using the  
15 Wisconsin Sequence Analysis Package Version 8-UNIX  
(Genetics Computer Group, Madison WI) and the GRAIL  
Sequence Analysis, Gene Assembly and Sequence  
Comparison System v. 1.2 (Informatics Group, Oak Ridge  
TN). Protein site motifs were identified using Motif  
20 (Genetics Computer Group, Madison WI).

Sources of Herpesvirus Gene Sequence Comparisons:  
Complete genomic sequences of three gammaherpesviruses  
were available: Epstein-Barr virus (EBV), a  
25 herpesvirus of humans [4]; herpesvirus saimiri (HVS),  
a herpesvirus of the New World monkey *Saimiri sciureus*  
[1]; and equine herpesvirus 2 (EHV2 [49]). Additional  
thymidine kinase gene sequences were obtained for  
alcelaphine herpesvirus 1 (AHV1 [22]) and bovine  
30 herpesvirus 4 (BHV4 [31]). Sequences for the major  
capsid protein genes of human herpesvirus 6B and human  
herpesvirus 7 (HHV7) were from Mukai et al. [34]. The  
sources of all other sequences used are listed  
previously in McGeoch and Cook [31] and McGeoch et al.  
35 [32].

Phylogenetic Inference: Predicted amino acid sequences used for tree construction were based on previous experience with herpesviral phylogenetic analyses [31]. Alignments of homologous sets of amino acid sequences were made with the AMPS [5] and Pileup [16] programs. Regions of alignments that showed extreme divergence with marked length heterogeneity, typically terminal sections, were excised. Generally, positions in alignments that contained inserted gaps in one or more sequences were removed before use for tree construction. Phylogenetic inference programs were from the Phylip set, version 3.5c [14] and from the GCG set [16]. Trees were built with the maximum parsimony (MP), neighbor joining (NJ) methods. For the NJ method, which utilizes estimates of pairwise distances between sequences, distances were estimated as mean numbers of substitution events per site with ProtDist using the PAM 250 substitution probability matrix of Schwartz & Dayhoff [46]. Bootstrap analysis [15] was carried out for MP and NJ trees, with 100 sub-replicates of each alignment, and consensus trees obtained with the program Consense. In addition the program Protml was used to infer trees by the maximum likelihood (ML) method. Protml was obtained from J. Adachi, Department of Statistical Science, The Graduate University for Advanced Study, Tokyo 106, Japan. Because of computational constraints, Protml was used only with the 4-species CS1 alignment.

Clamped Homogeneous Electric Field (CHEF) Gel Electrophoresis: Agarose plugs were prepared by resuspending BCBL-1 cells in 1% LMP agarose (Biorad, Hercules CA) and 0.9% NaCl at 42°C to a final concentration of  $2.5 \times 10^7$  cells/ml. Solidified agarose plugs were transferred into lysis buffer (0.5M EDTA pH 8.0, 1% sarcosyl, proteinase K at 1 mg/ml

final concentration) and incubated for 24 hours. Approximately  $10^7$  BCBL-1 cells were loaded in each lane. Gels were run at a gradient of 6.0 V/cm with a run time of 28 h 28 min. on a CHEF Mapper XA pulsed field gel electrophoresis apparatus (Biorad, Hercules CA), Southern blotted and hybridized to KS627Bam, KS330Bam and an EBV terminal repeat sequence [40].

TPA Induction of Genome Replication: Late log phase BCBL-1 cells ( $5 \times 10^5$  cells per ml) were incubated with varying amounts of 12-O-tetradecanoylphorbol-13-acetate (TPA, Sigma Chemical Co., St. Louis MO) for 48 h, cells were then harvested and washed with phosphate-buffered saline (PBS) and DNA was isolated by chloroform-phenol extraction. DNA concentrations were determined by UV absorbance; 5  $\mu$ g of whole cell DNA was quantitatively dot blot hybridized in triplicate (Manifold I, Schleicher and Schuell, Keene NH). KS631Bam, EBV terminal repeat and beta-actin sequences were random-primer labeled with  $^{32}$ P [13]. Specific hybridization was quantitated on a Molecular Dynamics PhosphorImager 425E.

Cell Cultures and Transmission Studies: Cells were maintained at  $5 \times 10^5$  cells per ml in RPMI 1640 with 20% fetal calf serum (FCS, Gibco-BRL, Gaithersburg MD) and periodically examined for continued KSHV infection by PCR and dot hybridization. The T cell line Molt-3 (a gift from Dr. Jodi Black, Centers for Disease Control and Prevention), Raji cells (American Type Culture Collection, Rockville MD) and RCC-1 cells were cultured in RPMI 1640 with 10% FCS. Owl monkey kidney cells (American Type Culture Collection, Rockville MD) were cultured in MEM with 10% FCS and 1% nonessential amino acids (Gibco-BRL, Gaithersburg MD).

To produce the RCC-1 cell line,  $2 \times 10^6$  Raji cells were cultivated with  $1.4 \times 10^6$  BCBL-1 cells in the presence of 20 ng/ml TPA for 2 days in chambers separated by Falcon 0.45  $\mu$ g filter tissue culture inserts to prevent contamination of Raji with BCBL-1. Demonstration that RCC-1 was not contaminated with BCBL-1 was obtained by PCR typing of HLA-DR alleles [27] (Raji and RCC-1: DR $\beta$ 1\*0310, DR $\beta$ 3\*02; BCBL-1: DR $\beta$ 104,\*07, Dr $\beta$ 4\*01) and confirmed by flow cytometry to determine the presence (Raji, RCC1) or absence (BCBL-1) of EMA membrane antigen. Clonal sublines of RCC-1 were obtained by dilution in 96 well plates to 0.1 cells/well in RPMI 1640, 20% FCS and 30% T-STIM culture supplement (Collaborative Biomedical Products, Bedford MA). Subcultures were examined to ensure that each was derived from a single cluster of growing cells.

*In situ* hybridization was performed with a previously described 25 bp oligomer located in ORF26 which was 5' labeled with fluorescein (Operon, Alameda CA) and hybridized to cytospin preparations of BCBL-1, RCC-1 and Raji cells using the methods of Lungu et al. [29]. Slides were both directly visualized by UV microscopy and by incubating slides with anti-fluorescein-alkaline phosphatase (AP)-conjugated antibody (Boehringer-Mannheim, Indianapolis IN), allowing immunohistochemical detection of bound probe. Positive control hybridization was performed using a 26 bp TET-labeled EBV DNA polymerase gene oligomer (Applied Biosystems, Alameda CA) which was visualized by UV microscopy only and negative control hybridization was performed using a 25 bp 5' fluorescein-labeled HSV1  $\alpha$ 47 gene oligomer (Operon, Alameda CA) which was visualized in a similar manner as the KSHV ORF26

probe. All nuclei of BCBL-1, RCC-1 and Raji appropriately stained with the EBV hybridization probe whereas no specific staining of the cells occurred after hybridization with the HSV1 probe.

5

The remaining suspension cell lines used in transmission experiments were pelleted, and resuspended in 5 ml of 0.22 or 0.45  $\mu$  filtered BCBL-1 tissue culture supernatant for 16 h. BCBL-1 supernatants were either from unstimulated cultures or from cultures stimulated with 20 ng/ml TPA. No difference in transmission to recipient cell lines was noted using various filtration or stimulation conditions. Fetal cord blood lymphocytes (FCBL) were obtained from heparinized fresh post-partum umbilical cord blood after separation on Ficoll-Paque (Pharmacia LKB, Uppsala Sweden) gradients and cultured in RPMI 1640 with 10% fetal calf serum. Adherent recipient cells were washed with sterile Hank's Buffered Salt Solution (HBSS, Gibco-BRL, Gaithersburg MD) and overlaid with 5 ml of BCBL-1 media supernatant. After incubation with BCBL-1 media supernatant, cells were washed three times with sterile HBSS, and suspended in fresh media. Cells were subsequently rewashed three times every other day for six days and grown for at least two weeks prior to DNA extraction and testing. PCR to detect KSHV infection was performed using nested and un-nested primers from ORF 26 and ORF 25 as previously described [10, 35].

35

Indirect Immunofluorescence Assay: AIDS-KS sera were obtained from ongoing cohort studies (provided by Drs. Scott Holmberg, Thomas Spira and Harold Jaffe, Centers for Disease Control, and Prevention, and Isaac Weisfuse, New York City Department of Health).

Sera from AIDS-KS patients were drawn between 1 and 31 months after initial KS diagnosis, sera from intravenous drug user and homosexual/bisexual controls were drawn after non-KS AIDS diagnosis, and 5 sera from HIV-infected hemophilic controls were drawn at various times after HIV infection. Immunofluorescence assays were performed using an equal volume mixture of goat anti-human IgG-FITC conjugate (Molecular Probes, Eugene OR) and goat 10 anti-human IgM-FITC conjugate (Sigma Chemical Co., St. Louis MO) diluted 1:100 and serial dilutions of patient sera. End-point titers were read blindly and specific immunoglobulin binding was assessed by the presence or absence of a specular fluorescence 15 pattern in the nuclei of the plated cells. To adsorb cross-reacting antibodies, 20  $\mu$ l serum diluted 1:10 in phosphate-buffer saline (PBS), pH 7.4, were adsorbed with  $1-3 \times 10^7$  paraformaldehyde-fixed P3H3 cells for 4-10 h at 25° C and removed by 20 low speed centrifugation. P3H3 were induced prior to fixation with 20 ng/ml TPA for 48 h, fixed with 1% paraformaldehyde in PBS for 2 h at 4° C, and washed three times in PBS prior to adsorption.

## 25 RESULTS

### Sequence Analysis of a 20.7 kb KSHV DNA Sequence:

To demonstrate that KS330Bam and KS631Bam are genomic fragments from a new and previously 30 uncharacterized herpesvirus, a lambda phage clone (KS5) derived from an AIDS-KS genomic DNA library was identified by hybridization to the KS330Bam sequence. The KS5 insert was subcloned after NotI/BamHI digestion into five subfragments and both 35 strands of each fragment were sequenced by primer walking or nested deletion with a 7-fold average



redundancy. The KS5 sequence is 20,705 bp in length and has a G+C content of 54.0%. The observed/expected CpG dinucleotide ratio is 0.92 indicating no overall CpG suppression in this region.

Open reading frame (ORF) analysis identified 15 complete ORFs with coding regions ranging from 231 bp to 4128 bp in length, and two incomplete ORFs at the termini of the KS5 clone which were 135 and 552 bp in length (Figure 12). The coding probability of each ORF was analyzed using GRAIL 2 and CodonPreference which identified 17 regions having excellent to good protein coding probabilities. Each region is within an ORF encoding a homolog to a known herpesvirus gene with the exception of one ORF located at the genome position corresponding to ORF28 in herpesvirus saimiri (HVS). Codon preference values for all of the ORFs were higher across predicted ORFs than in non-coding regions when using a codon table composed of KS5 homologs to the conserved herpesvirus major capsid (MCP), glycoprotein H (gH), thymidine kinase (TK), and the putative DNA packaging protein (ORF29a/ORF29b) genes.

The translated sequence of each ORF was used to search GenBank/EMBL databases with BLASTX and FastA algorithms [2, 38]. All of the putative KS5 ORFs, except one, have sequence and collinear positional homology to ORFs from gamma-2 herpesviruses, especially HVS and equine herpesvirus 2 (EHV2). Because of the high degree of collinearity and amino acid sequence similarity between KSHV and HVS, KSHV ORFs have been named according to their HVS

positional homologs (i.e. KSHV ORF25 is named after HVS ORF 25).

5 The KS5 sequence spans a region which includes three  
of the seven conserved herpesvirus gene blocks  
(Figure 14) [10]. ORFs present in these blocks  
include genes which encode herpesvirus virion  
structural proteins and enzymes involved in DNA  
10 metabolism and replication. Amino acid identities  
between KS5 ORFs and HVS ORFs range from 30% to 60%,  
with the conserved MCP ORF25 and ORF29b genes having  
the highest percentage amino acid identity to  
homologs in other gammaherpesviruses. KSHV ORF28,  
15 which has no detectable sequence homology to HVS or  
EBV genes, has positional homology to HVS ORF28 and  
EBV BDLF3. ORF28 lies at the junction of two gene  
blocks (Figure 14); these junctions tend to exhibit  
greater sequence divergence than intrablock regions  
among herpesviral genomes [17]. Two ORFs were  
20 identified with sequence homology to the putative  
spliced protein packaging genes of HVS  
(ORF29a/ORF29b) and herpes simplex virus type 1  
(UL15). The KS330Bam sequence is located within  
KSHV ORF26, whose HSV-1 counterpart, VP23, is a  
25 minor virion structural component.

For every KSHV homolog, the HVS amino acid  
similarity spans the entire gene product, with the  
exception of ORF21, the TK gene. The KSHV TK  
30 homolog contains a proline-rich domain at its amino  
terminus (nt 20343-19636; aa 1-236) that is not  
conserved in other herpesvirus TK sequences, while  
the carboxyl terminus (nt 19637-18601; aa 237-565)  
is highly similar to the corresponding regions of  
35 HVS, EHV2, and bovine herpesvirus 4 (BHV4) TK. A  
purine binding motif with a glycine-rich region

found in herpesviral TK genes, as well as other TK genes, is present in the KSHV TK homolog (GVMGVGKS; aa 260-267).

5 The KS5 translated amino acid sequences were searched against the PROSITE Dictionary of Protein Sites and Patterns (Dr. Amos Bairoch, University of Geneva, Switzerland) using the computer program Motifs. Four sequence motif matches were identified  
10 among KSHV hypothetical protein sequences. These matches included: (i) a cytochrome c family heme-binding motif in ORF33 (CVHCHG; aa 209-214) and ORF34 (CLLCHI; aa 257-261), (ii) an immunoglobulin and major histocompatibility complex protein signature in ORF25  
15 (FICQAKH; aa 1024-1030), (iii) a mitochondrial energy transfer protein motif in ORF26 (PDDITRMRV; aa 260-268), and (iv) the purine nucleotide binding site identified in ORF21. The purine binding motif is the only motif with obvious functional significance. A  
20 cytosine-specific methylase motif present in HVS ORF27 is not present in KSHV ORF27. This motif may play a role in the methylation of episomal DNA in cells persistently infected with HVS [1].

25 Phylogenetic Analysis of KSHV: Amino acid sequences translated from the KS5 sequence were aligned with corresponding sequences from other herpesviruses. On the basis of the level of conserved aligned residues and the low incidence of introduced gaps, the amino acid  
30 alignments for ORFs 21, 22, 23, 24, 25, 26, 29a, 29b, 31 and 34 were suitable for phylogenetic analyses.

To demonstrate the phylogenetic relationship of KSHV to other herpesviruses, a single-gene comparison was made  
35 for ORF25 (MCP) homologs from KS5 and twelve members of Herpesviridae (Figures 15A-15B). The thirteen available MCP amino acid sequences are large (1376 a.a. residues for the KSHV homolog) and alignment required only a low

level of gapping; however, the overall similarity between viruses is relatively low [33]. The MCP set gave stable trees with high bootstrap scores and assigned the KSHV homolog to the gamma-2 sublineage (genus *Rhadinovirus* ), containing HVS, EHV2 and BVH4 [20, 33, 43]. KSHV was most closely associated with HVS. Similar results were obtained for single-gene alignments of TK and UL15/ORF29 sets but with lower bootstrap scores so that among gamma-2 herpesvirus members branching orders for EHV2, HVS and KSHV were not resolved.

To determine the relative divergence between KSHV and other gammaherpesviruses, alignments for the nine genes listed above were concatenated to produce a combined gammaherpesvirus gene set (CS1) containing EBV, EHV2, HVS and KSHV amino acid sequences. The total length of CS1 was 4247 residues after removal of positions containing gaps introduced by the alignment process in one or more of the sequences. The CS1 alignment was analyzed by the ML method, giving the tree shown in Figure 15B and by the MP and NJ methods used with the aligned herpesvirus MCP sequences. All three methods identified KSHV and HVS as sister groups, confirming that KSHV belongs in the gamma-2 sublineage with HVS as its closest known relative. It was previously estimated that divergence of the HVS and EHV2 lineages may have been contemporary with divergence of the primate and ungulate host lineages [33]. The results for the CS1 set suggest that HVS and KSHV represent a lineage of primate herpesviruses and, based on the distance between KSHV and HVS relative to the position of EHV2, divergence between HVS and KSHV lines is ancient.

#### 35 Genomic Studies of KSHV:

CHEF electrophoresis performed on BCBL-1 cells embedded in agarose plugs demonstrated the presence of a nonintegrated KSHV genome as well as a high molecular weight species (Figures 16A-16B). KS631Bam (Figure 16A)

and KS330Bam specifically hybridized to a single CHEF gel band comigrating with 270 kilobase (kb) linear DNA standards. The majority of hybridizing DNA was present in a diffuse band at the well origin; a low intensity high molecular weight (HMW) band was also present immediately below the origin (Figure 16A. arrow). The same filter was stripped and probed with an EBV terminal repeat sequence [40] yielding a 150-160 kb band (Figure 16B) corresponding to linear EBV DNA [24]. The HMW EBV band may correspond to either circular or concatemeric EBV DNA [24].

The phorbol ester TPA induces replication-competent EBV to enter a lytic replication cycle [49]. To determine if TPA induces replication of KSHV and EBV in BCBL-1 cells, these cells were incubated with varying concentrations of TPA for 48 h (Figure 17). Maximum stimulation of EBV occurred at 20 ng/ml TPA which resulted in an eight-fold increase in hybridizing EBV genome. Only a 1.3-1.4 fold increase in KSHV genome abundance occurred after 20-80 ng/ml TPA incubation for 48 h.

#### Transmission Studies:

Prior to determining that the agent was likely to be a member of Herpesviridae by sequence analysis, BCBL-1 cells were cultured with Raji cells, a nonlytic EBV transformed B cell line, in chambers separated by a 0.45  $\mu$  tissue culture filter. Recipient Raji cells generally demonstrated rapid cytolysis suggesting transmission of a cytotoxic component from the BCBL-1 cell line. One Raji line cultured in 10 ng/ml TPA for 2 days, underwent an initial period of cytolysis before recovery and resumption of logarithmic growth. This cell line (RCC-1) is a monoculture derived from Raji uncontaminated by BCBL-1 as determined by PCR amplification of HLA-DR sequences.

RCC-1 has remained positive for the KS330<sub>233</sub> PCR product for >6 months in continuous culture (approximately 70 passages), but KSHV was not detectable by dot or Southern hybridization at any time. In situ hybridization, however, with a 25 bp KSHV ORF26-derived oligomer was used to demonstrate persistent localization of KSHV DNA to RCC-1 nuclei. As indicated in Figures 18A-18C, nuclei of BCBL-1 and RCC-1 (from passage ~65) cells had detectable hybridization with the ORF26 oligomer, whereas no specific hybridization occurred with parental Raji cells (Figure 18B). KSHV sequences were detectable in 65% of BCBL-1 and 2.6% of RCC-1 cells under these conditions. In addition, forty-five monoclonal cultures were subcultured by serial dilution from RCC-1 at passage 50, of which eight (18%) clones were PCR positive by KS330<sub>233</sub>. While PCR detection using unnested KS330<sub>233</sub> primer pairs was lost by passage 15 in each of the clonal cultures, persistent KSHV genome was detected in 5 clones using two more sensitive nonoverlapping nested PCR primer sets [33] suggesting that KSHV genome is lost over time in RCC-1 and its clones.

Low but persistent levels of KS330<sub>233</sub> PCR positivity were found for one of four Raji, one of four Bjab, two of three Molt-3, one of one owl monkey kidney cell lines and three of eight human fetal cord blood lymphocyte (FCBL) cultures after inoculation with 0.2-0.45  $\mu$  filtered BCBL-1 supernatants. Among the PCR positive cultures, PCR detectable genome was lost after 2-6 weeks and multiple washings. Five FCBL cultures developed cell clusters characteristic of EBV immortalized lymphocytes and were positive for EBV by PCR using EBER primers [23]; three of these cultures were also initially KS330<sub>233</sub> positive. None of the recipient cell lines had detectable KSHV genome by dot blot hybridization.

Serologic Studies:

Indirect immunofluorescence antibody assays (IFA) were used to assess the presence of specific antibodies against the KSHV- and EBV-infected cell line HBL-6 in the sera from AIDS-KS patients and control patients with HIV infection or AIDS. HBL-6 was substituted for BCBL-1 for reasons of convenience; preliminary studies showed no significant differences in IFA results between HBL-6 and BCBL-1. HBL-6 have diffuse immunofluorescent cell staining with most KS patient and control unadsorbed sera suggesting nonspecific antibody binding (Figures 19A-19D). After adsorption with paraformaldehyde-fixed, TPA-induced P3H3 (an EBV producer subline of P3J-HR1, a gift of Dr. George Miller) to remove cross-reacting antibodies against EBV and lymphocyte antigens, patient sera generally showed specular nuclear staining at high titers while this staining pattern was absent from control patient sera (Figures 19B and 19D). Staining was localized primarily to the nucleus but weak cytoplasmic staining was also present at low sera dilutions.

With unadsorbed sera, the initial endpoint geometric mean titers (GMT) against HBL-6 cell antigens for the sera from AIDS-KS patients (GMT=1:1153, range: 1:150 to 1:12,150) were higher than for sera from control, non-KS patients (GMT=1:342, range 1:50 to 1:12,150;  $p=0.04$ ) (Figure 13). While AIDS-KS patients and HIV-infected gay/bisexual and intravenous drug user control patients had similar endpoint titers to HBL-6 antigens (GMT=1:1265 and GMT=1:1578, respectively), hemophilic AIDS patient titers were lower (GMT=1:104). Both case and control patient groups had elevated IFA titers against the EBV infected cell line P3H3.

The difference in endpoint GMT between case and control titers against HBL-6 antigens increased after adsorption with P3H3. After adsorption, case GMT declined to 1:780 and control GMT declined to 1:81 ( $p=0.00009$ ). Similar

SUB  
B10

5 results were obtained by using BCBL-1 instead of BHL-6 cells, by pre-adsorbing with EBV-infected nonproducer Raji cells instead of P3H3 and by using sera from a homosexual male KS patient without HIV infection, in complete remission for KS for 9 months (BHL-6 titer 1:450, P3H3 titer 1:150). Paired sera taken 8-14 months prior to KS onset and after KS onset were available for three KS patients: KS patients 8 and 13 had eight-fold rises and patient 8 had a three-fold fall in P3H3-adsorbed BCBL-1 titers from pre-onset sera to post-KS sera.

DISCUSSION

15 These studies demonstrate that specific DNA sequences found in KS lesions by representational difference analysis belong to a newly identified human herpesvirus. The current studies define this agent as a human gamma-2 herpesvirus that can be continuously cultured in naturally-transformed, EBV-coinfected lymphocytes from AIDS-related body-cavity based lymphomas.

25 Sequence analysis of the KS5 lambda phage insert provides clear evidence that the KS330Bam sequence is part of a larger herpesvirus genome. KS5 has a 54.0% G+C content which is considerably higher than the corresponding HVS region (34.3% G+C). While there is no CpG dinucleotide suppression in the KS5 sequence, the corresponding HVS region has a 0.33 expected:observed CpG dinucleotide ratio [1]. The CpG dinucleotide frequency in herpesviruses varies from global CpG suppression among gammaherpesviruses to local CpG suppression in the beta herpesviruses, which may result from deamination of 5'-methylcytosine residues at CpG sites resulting in TpG substitutions [21]. CpG suppression among herpesviruses [21, 30, 44] has been hypothesized to reflect co-replication of latent genome in actively dividing host cells, but it is unknown whether or not KSHV is primarily maintained by a lytic replication cycle in vivo.

000290 04740360



5 The 20,705 bp KS5 fragment has 17 protein-coding regions, 15 of which are complete ORFs with appropriately located TATA and polyadenylation signals, and two incomplete ORFs located at the phage insert  
10 termini. Sixteen of these ORFs correspond by sequence and collinear positional homology to 15 previously identified herpesviral genes including the highly conserved spliced gene. The conserved positional and sequence homology for KSHV genes in this region are  
15 consistent with the possibility that the biological behavior of the virus is similar to that of other gammaherpesviruses. For example, identification of a thymidine kinase-like gene on KS5 implies that the agent is potentially susceptible to TK-activated DNA  
20 polymerase inhibitors and like other herpesviruses possesses viral genes involved in nucleotide metabolism and DNA replication [41]. The presence of major capsid protein and glycoprotein H gene homologs suggest that replication competent virus would produce a capsid structure similar to other herpesviruses.

Phylogenetic analyses of molecular sequences show that KSHV belongs to the gamma-2 sublineage of the Gammaherpesvirinae subfamily, and is thus the first  
25 human gamma-2 herpesvirus identified. Its closest known relative based on available sequence comparisons is HVS, a squirrel monkey gamma-2 herpesvirus that causes fulminant polyclonal T cell lymphoproliferative disorders in some New World monkey species. Data for  
30 the gamma-2 sublineage are sparse: only three viruses (KSHV, HVS and EHV2) can at present be placed on the phylogenetic tree with precision (the sublineage also contains murine herpesvirus 68 and BHV4 [33]). Given the limitation in resolution imposed by this thin  
35 background, KSHV and HVS appear to represent a lineage of primate gamma-2 viruses. Previously, McGeoch et al. [33] proposed that lines of gamma-2 herpesviruses may have originated by cospeciation with the ancestors of their host species. Extrapolation of this view to KSHV

and HVS suggests that these viruses diverged at an ancient time, possibly contemporaneously with the divergence of the Old World and New World primate host lineages. Gammaherpesviruses are distinguished as a subfamily by their lymphotropism [41] and this grouping is supported by phylogenetic analysis based on sequence data [33]. The biologic behavior of KSHV is consistent with its phylogenetic designation in that KSHV can be found in in vitro lymphocyte cultures and in in vivo samples of lymphocytes [3].

This band appears to be a linear form of the genome because other "high molecular weight" bands are present for both EBV and KSHV in BCBL-1 which may represent circular forms of their genomes. The linear form of the EBV genome, associated with replicating and packaged DNA [41] migrates substantially faster than the closed circular form associated with latent viral replication [24]. While the 270 kb band appears to be a linear form, it is also consistent with a replicating dimer plasmid since the genome size of HVS is approximately 135 kb. The true size of the genome may only be resolved by ongoing mapping and sequencing studies.

Replication deficient EBV mutants are common among EBV strains passaged through prolonged tissue culture [23]. The EBV strain infecting Raji, for example, is an BALF-2 deficient mutant [19]; virus replication is not inducible with TPA and its genome is maintained only as a latent circular form [23, 33]. The EBV strain coinfecting BCBL-1 does not appear to be replication deficient because TPA induces eight-fold increases in DNA content and has an apparent linear form on CHEF electrophoresis. KSHV replication, however, is only marginally induced by comparable TPA treatment indicating either insensitivity to TPA induction or that the genome has undergone loss of genetic elements required for TPA induction. Additional experiments, however, indicate that KSHV DNA can be pelleted by high

speed centrifugation of filtered organelle-free, DNase I-protected BCBL-1 cell extracts, which is consistent with KSHV encapsidation.

5      Transmission of KSHV DNA from BCBL-1 to a variety of recipient cell lines is possible and KSHV DNA can be maintained at low levels in recipient cells for up to 70 passages. However, detection of virus genome in recipient cell lines by PCR may be due to physical association of KSHV DNA fragments rather than true infection. This appears to be unlikely given evidence for specific nuclear localization of the ORF26 sequence in RCC-1. If transmission of infectious virus from BCBL-1 occurs, it is apparent that the viral genome declines in abundance with subsequent passages of recipient cells. This is consistent with studies of spindle cell lines derived from KS lesions. Spindle cell cultures generally have PCR detectable KSHV genome when first explanted, but rapidly lose viral genome after initial passages and established spindle cell cultures generally do not have detectable KSHV sequences [3].

25      Infections with the human herpesviruses are generally ubiquitous in that nearly all humans are infected by early adulthood with six of the seven previously identified human herpesviruses [42]. Universal infection with EBV, for example, is the primary reason for the difficulty in clearly establishing a causal role for this virus in EBV-associated human tumors. The serologic studies identified nuclear antigen in BCBL-1 and BHL-6 which is recognized by sera from AIDS-KS patients but generally not by sera from control AIDS patients without KS after removal of EBV-reactive antibodies. These data are consistent with PCR studies of KS and control patient lymphocytes suggesting that KSHV is not ubiquitous among adult humans, but is specifically associated with persons who develop Kaposi's sarcoma. In this respect, it appears to be

SUB  
B11

005250 02F20500

epidemiologically similar to HSV2 rather than the other known human herpesviruses. An alternative possibility is that elevated IFA titers against BCBL-1 reflect disease status rather than infection with the virus.

REFERENCES

- 5 1. Albrecht, J.-C., J. Nicholas, D. Biller, K. R. Cameron, B. Biesinger, C. Newamn, S. Wittmann, M. A. Craxton, H. Coleman, B. Fleckenstein, and R. W. Honess. 1992. Primary structure of the Herpesvirus saimiri genome. *J Virol.* 66:5047-5058.
- 10 2. Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J Mol Biol.* 215:403-410.
- 15 3. Ambroziak, J. A., D. J. Blackbourn, B. G. Herndier, R. G. Glogau, J. H. Gullett, A. R. McDonald, E. T. Lennette, and J. A. Levy. 1995. Herpes-like sequences in HIV-infected and uninfected Kaposi's sarcoma patients. *Science.* 268:582-583.
- 20 4. Baer, R., A. T. Bankier, P. L. Biggin, P. L. Deininger, P. J. Farrell, T. J. Gibson, G. Hatfull, G. S. Hudson, S. C. Satchwell, C. Séguin, P. S. Tuffnell, and B. G. Barrell. 1984. DNA sequence and expression of the B95-8 Epstein-Barr virus genome. *Nature.* 310:207-211.
- 25 5. Barton, G. J., and M. J. E. Sternberg. 1987. A strategy for the rapid multiple alignment of protein sequences. Confidence levels from tertiary structure comparisons. *J Mol Biol.* 198:327-37.
- 30 6. Beral, V., T. A. Peterman, R. L. Berkelman, and H. W. Jaffe. 1990. Kaposi's sarcoma among persons with AIDS: a sexually transmitted infection? *Lancet.* 335:123-128.
- 35

006290" 62F20900

- 5
7. Boshoff, C., D. Whitby, T. Hatziionnou, C. Fisher, J. van der Walt, A. Hatzakis, R. Weiss, and T. Schulz. 1995. Kaposi's sarcoma-associated herpesvirus in HIV-negative Kaposi's sarcoma. *Lancet*. 345:1043-44.
- 10
8. Cesarman, E., Y. Chang, P. S. Moore, J. W. Said, and D. M. Knowles. 1995. Kaposi's sarcoma-associated herpesvirus-like DNA sequences are present in AIDS-related body cavity based lymphomas. *New England J Med*. 332:1186-1191.
- 15
9. Chang, Y., E. Cesarman, M. S. Pessin, F. Lee, J. Culpepper, D. M. Knowles, and P. S. Moore. 1994. Identification of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. *Science*. 265:1865-69.
- 20
10. Chee, M. S., S. B. Bankier, C. M. Bohni, R. C. Brown, T. Horsnell, C. A. Hutchison, T. Kouzarides, J. A. Martignetti, E. Preddie, S. C. Satchwell, P. Tomlinson, K. M. Weston, and B. G. Barrell. 1990. Analysis of the protein coding content of the sequence of cytomegalovirus strain AD169. *Curr Top Microbiol Immunol*. 154:125-69.
- 25
- 30
11. Collandre, H., S. Ferris, O. Grau, L. Montagnier, and A. Blanchard. 1995. Kaposi's sarcoma and new herpesvirus. *Lancet*. 345:1043.
- 35
12. Dupin, N., M. Grandadam, V. Calvez, I. Gorin, J. T. Aubin, S. Harvard, F. Lamy, M. Leibowitch, J. M. Huraux, J. P. Escande, and H. Agut. 1995. Herpesvirus-like DNA in patients with Mediterranean Kaposi's sarcoma. *Lancet*. 345:761-2.

005250\*62720960

- 5
- 10
- 15
- 20
- 25
- 30
- 35
13. Feinberg, A. P., and B. Vogelstein. 1983. A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem.* 132:6.
  14. Felsenstein, J. 1989. PHYLIP-phylogeny inference package (ver 3.2). *Cladistics.* 5:164-6.
  15. Felsenstein, J. 1988. Phylogenies from molecular sequences: inferences and reliability. *Annual Rev Microbiol.* 22:521-65.
  16. Genetics Computer Group. 1994. Program manual for the GCG package, version 8, Madison, Wisconsin.
  17. Gompels, U. A., J. Nicholas, G. Lawrence, M. Jones, B. J. Thomson, M. E. D. Martin, S. Efstathiou, M. Craxton, and H. A. Macaulay. 1995. The DNA sequence of human herpesvirus-6: Structure, coding content and genome evolution. *Virology.* 209:29-51.
  18. Hatfull, G., A. T. Bankier, B. G. Barrell, and P. J. Farrell. 1988. Sequence analysis of Raji Epstein-Barr virus DNA. *Virol.* 164:334-40.
  19. Holmberg, S. D. 1990. Possible cofactors for the development of AIDS-related neoplasms. *Cancer Detection and Prevention.* 14:331-336.
  20. Honess, R. W. 1984. Herpes simplex and 'the herpes complex': diverse observations and a unifying hypothesis. *J Gen Virol.* 65:2077-2107.

006230-6272990

- 5 21. Honess, R. W., U. A. Gompels, B. G. Barrell, M. Craxton, K. R. Cameron, R. Staden, Y.-N. Chang, and G. S. Hayward. 1989. Deviations from expected frequencies of CpG dinucleotides in herpesvirus DNAs may be diagnostic of differences in the states of their latent genomes. *J Gen Virol.* 70:837-55.
- 10 22. Hsu, D., L. M. Shih, and Y. C. Zee. 1990. Nucleotide sequence of a 3.5 nucleotide fragment of malignant catarrhal fever virus strain WC11. *Arch Virol.* 113:53-60.
- 15 23. Kieff, E., and D. Liebowitz. 1990. Epstein-Barr virus and its replication, p. 1889-1920. In B. N. Fields and D. M. Knipe (ed.), *Virology*, vol. 2. Raven Press, New York.
- 20 24. Kolman, J. L., C. J. Kolman, and G. Miller. 1992. Marked variation in the size of genomic plasmids among members of the family of related Epstein-Barr viruses. *Proc Natl Acad Sci, USA.* 89:7772-7776.
- 25 25. Lebbé, C., P. de Crémoux, M. Rybojad, C. Costa da Cunha, P. Morel, and F. Calvo. 1995. Kaposi's sarcoma and new herpesvirus. *Lancet.* 345:1180.
- 30 26. Lin, J. C., S. C. Lin, B. K. De, W. P. Chan, and B. L. Evatt. 1993. Precision of genotyping of Epstein-Barr virus by polymerase chain reaction using three gene loci (EBNA-2, EBNA-3C and EBER): predominance of type A virus associated with Hodgkin's disease. *Blood.* 81:3372-81.
- 35 27. Liu, Z., S. Yu-Kai, Y.-P. Xi, P. Harris, and N. Suciú-Foca. 1992. T cell recognition of

EPSTEIN-BARR VIRUS



self-human histocompatibility . leukocyte antigens (HLA)-DR peptides in the context of syngeneic HLA-DR molecules. *J Exp Med.* 175:1663-8.

- 5
28. Lomonte, P., M. Bublot, P-P. Pastoret, and E. Thiry. 1992. Location and characterization of the bovine herpesvirus type 4 thymidine kinase gene; comparison with thymidine kinase of other herpesviruses. *Arch. Virol.* 127:327-337.
- 10
29. Martin, M. E. D., J. Nicholas, B. J. Thomson, C. Newman, and R. W. Honess. 1991. Identification of a transactivating function mapping to the putative immediate-early locus of human herpesvirus 6. *J Virol.* 65:5381-5390.
- 15
- 20
30. McGeoch, D. J., and S. Cook. 1994. Molecular phylogeny of the Alphaherpesvirinae subfamily and a proposed evolutionary timescale. *J Mol Biol.* 238:9-22.
- 25
31. McGeoch, D. J., S. Cook, A. Dolan, F. E. Jamieson, and E. A. R. Telford. 1995. Molecular phylogeny and evolutionary timescale for the family of mammalian herpesviruses. *J Molec Biol.* 247:443-58.
- 30
32. Miller, G. 1990. Epstein-Barr virus: Biology, pathogenesis and medical aspects, p. 1921-1957. In B. N. Fields and D. M. Knipe (ed.), *Virology*, 2nd ed, vol. 2. Raven Press, New York.
- 35
33. Moore, P. S., and Y. Chang. 1995. Detection of herpesvirus-like DNA sequences in Kaposi's sarcoma lesions from persons with and without

HIV infection. *New England J Med.* 332:1181-1185.

- 5           34. Mukai, T., Y. Isegawa, and K. Yamanishi. 1995. Identification of the major capsid protein gene of human herpesvirus 7. *Virus Res.* 37:55-62.
- 10           35. Oettle, A. G. 1962. Geographic and racial differences in the frequency of Kaposi's sarcoma as evidence of environmental or genetic causes, vol. 18. Symposium on Kaposi's sarcoma: *Unio Internationalis Contra Cancrum*, Karger, Basel.
- 15
36. Pearson, W. R., and D. J. Lipman. 1988. Improved tools for biological sequence analysis. *Proc Natl Acad Sci, USA.* 85:2444-8.
- 20
37. Peterman, T. A., H. W. Jaffe, A. E. Friedman-Kien, and R. A. Weiss. 1991. The aetiology of Kaposi's sarcoma, p. 23-37, *Cancer, HIV, and AIDS*, vol. 10. Imperial Cancer Research Fund. London.
- 25
38. Raab-Traub, N., and K. Flynn. 1986. The structure of the termini of the Epstein-Barr virus as a marker of clonal cellular proliferation. *Cell.* 47:883-889.
- 30
39. Roizman, B. 1993. The family Herpesviridae, p. 1-9. In B. Roizman and R. J. Whitley and C. Lopez (ed.), *The Human Herpeviruses*. Raven Press, Ltd., New York.
- 35
40. Roizman, B. 1995. New viral footprints in Kaposi's sarcoma. *N Engl J Med.* 332:1227-1228.

- 5
41. Roizman, B., R. C. Desrosiers, B. Fleckenstein, C. López, A. C. Minson, and M. J. Studdert. 1992. The family Herpesviridae: an update. *Arch Virol.* 123:425-449.
- 10
42. Sandford, G. R., K. Ho, and W. H. Burns. 1993. Characterization of the major locus of immediate-early genes of rat cytomegalovirus. *J Virol.* 67:4093-4103.
- 15
43. Schalling, M., M. Ekman, E. E. Kaaya, A. Linde, and P. Bieberfeld. 1995. A role for a new herpesvirus (KSHV) in different forms of Kaposi's sarcoma. *Nature Med.* 1:707-8.
- 20
44. Schwartz, R. M., and M. O. Dayhoff. 1978. Matrices for detecting distant relationships, p. 353-8. In M. O. Dayhoff (ed.), *Atlas of protein sequence and structure*, vol. 5, suppl 3. National Biomedical Research Foundation, Washington.
- 25
45. Su, I.-J., Y.-S. Hsu, Y.-C. Chang, and I.-W. Wang. 1995. Herpesvirus-like DNA sequence in Kaposi's sarcoma from AIDS and non-AIDS patients in Taiwan. *Lancet.* 345:722-23.
- 30
46. Telford, E. A. R., M. S. Watson, H. C. Aird, J. Perry, and A. J. Davison. 1995. The DNA sequence of equine herpesvirus 2. *J Molec Biol.* 249:520-8.
- 35
47. zur Hausen, H., F. J. O'Neill, and U. K. Freese. 1978. Persisting oncogenic herpesvirus induced by the tumor promoter TPA. *Nature.* 272:373-375.

EXPERIMENTAL DETAILS SECTION III:

5 KS Patient Enrollment: Cases and controls were selected from ongoing cohort studies based on the availability of clinical information and appropriate PBMC samples. 21 homosexual or bisexual men with AIDS who developed KS during their participation in prospective cohort studies were identified [14-16]. Fourteen of these patients had paired PBMC samples collected after KS diagnosis (median 10 +4 months) and at least four months prior to KS diagnosis (median -13 months), while the remaining 7 had paired PBMC taken at the study visit immediately prior to KS diagnosis (median -3 months) and at entry into their cohort study (median -51 months prior to KS 15 diagnosis).

20 Hemophilic and Homosexual/Bisexual Male AIDS Patient Control Enrollment: Two control groups of AIDS patients were examined: 23 homosexual/bisexual men with AIDS followed until death who did not develop KS ("high risk" control group) from the Multicenter AIDS Cohort Study [16]), and 19 hemophilic men ("low risk" control group) enrolled from joint projects of the National Hemophilia Foundation and the Centers for Disease Control and 25 Prevention. Of the 19 hemophilic controls with available follow-up information, none are known to have developed KS and <2% of hemophilic AIDS patients historically develop KS [2]. For homosexual/bisexual AIDS control patients who did not develop KS, paired 30 PBMC specimens were available at entry into their cohort study (median -35 months prior to AIDS onset) and at the study visit immediately prior to nonKS AIDS diagnosis (median BHL-6 months prior to AIDS onset).

35 DNA Extraction and Analyses: DNA from  $10^6$ - $10^7$  PBMC in each specimen was extracted and quantitated by spectrophotometry. Samples were prepared in physically isolated laboratories from the laboratory where polymerase chain reaction (PCR) analyses were performed.

000000-000000

SUB  
B12

All samples were tested for amplifiability using primers specific for either the HLA-DQ locus (GH26/GH27) or b-globin [18]. PCR detection of KSHV DNA was performed as previously described [7] with the following nested primer sets: No. 1 outer 5'-AGCACTCGCAGGGCAGTACG-3', 5'-GACTCTTCGCTGATGAACTGG-3'; No. 1 inner 5'-TCCGTGTTGTCTACGTCCAG-3', 5'-AGCCGAAAGGATTCCACCAT-3'; No. 2 outer 5'-AGGCAACGTCAGATGTGAC-3', 5'-GAAATTACCCACGAGATCGC-3'; No. 2 inner 5'-CATGGGAGTACATTGTCAGGACCTC-3', 5'-GGAATTATCTCGCAGGTTGCC-3'; No. 3 outer 5'-GGCGACATTCATCAACCTCAGGG-3', 5'-ATATCATCCTGTGCGTTCACGAC-3'; No. 3 inner 5'-CATGGGAGTACATTGTCAGGACCTC-3', 5'-GGAATTATCTCGCAGGTTGCC-3'. The outer primer set was amplified for 35 cycles at 94° C for 30 seconds, 60° C for 1 minute and 72° C for 1 minute with a 5 minute final extension cycle at 72° C. One to three ml of the PCR product was added to the inner PCR reaction mixture and amplified for 25 additional cycles with a 5 minute final extension cycle. Primary determination of sample positivity was made with primer set No. 1 and confirmed with either primer sets 2 or 3 which amplify nonoverlapping regions of the KSHV hypothetical major capsid gene. Sampling two portions of the KSHV genome decreased the likelihood of intraexperimental PCR contamination. These nested primer sets are 2-3 logs more sensitive for detecting KSHV sequences than the previously published KS330<sub>233</sub> primers [6] and are estimated to be able to detect <10 copies of KSHV genome under optimal conditions. Sample preparations were prealiquoted and amplified with alternating negative control samples without DNA to monitor and control possible contamination. All samples were tested in a blinded fashion and a determination of the positivity/negativity made before code breaking. Significance testing was performed with Mantel-Haenszel chi-squared estimates and exact confidence intervals using Epi-Info ver. 6 (USD Inc., Stone Mt. GA).

SECRET  
SUB  
B3

**RESULTS****KSHV Positivity of Case and Control PBMC Samples:**

5 Paired PBMC samples were available from each KS patient and homosexual/bisexual control patient; a single sample was available from each hemophilic control patient.

10 To determine the KSHV positivity rate for each group of AIDS patients, a single specimen from each participant taken closest to KS or other AIDS-defining illness ("second sample") was analyzed. Overall, 12 of 21 (57%) of PBMC specimens from KS patients taken from 6 months prior to KS diagnosis to 20 months after KS diagnosis were KSHV positive. There was no apparent difference in  
15 positivity rate between immediate pre-diagnosis and post-diagnosis visit specimens (4 of 7 (57%) vs. 8 of 14 (57%) respectively).

20 The number of KSHV positive control PBMC specimens from both homosexual/bisexual (second visit) and hemophilic patient controls was significantly lower. Only 2 of 19 (11%) hemophilic PBMC samples were positive (odds ratio 11.3, 95 % confidence interval 1.8 to 118) and only 2 of 23 (9%) PBMC samples from homosexual/bisexual men who  
25 did not develop KS were positive (odds ratio 14.0, 95% confidence interval 2.3 to 144). If all KS patient PBMC samples taken immediately prior to or after diagnosis were truly infected, the PCR assay was at least 57% sensitive in detecting KSHV infection among PBMC  
30 samples. No significant differences in CD4+ counts were found for KS patients and homosexual/bisexual patients without KS at the second sample evaluation (Kruskall-Wallis  $p=0.15$ ) (Figure 21). CD4+ counts from the single sample from hemophilic AIDS patients were higher than  
35 CD4+ counts from KS patients (Kruskall-Wallis  $p=0.004$ ), although both groups showed evidence of HIV-related immunosuppression.

Longitudinal Studies:

Paired specimens were available from all 21 KS patients and 23 homosexual/bisexual male AIDS control patients who did not develop KS. For the KS group, initial PBMC samples were taken four to 87 months (median 13 months) prior to the onset of KS. Initial PBMC samples from the control group were drawn 13 to 106 months (median 55 months) prior to onset of first nonKS AIDS-defining illness (1987 CDC surveillance definition). 11 of 21 (52%) of KS patients had detectable KSHV DNA in PBMC samples taken prior to KS onset compared to 2 of 19 (11%,  $p=0.005$ ) hemophilic control samples, and 1 (4%,  $p=0.0004$ ) and 2 (9%,  $p=0.002$ ) of 23 homosexual/bisexual control samples taken at the first and second visits respectively (Figures 20A-20B). The figure shows that 7 of the paired KS patient samples were positive at both visits, 5 KS patients and 2 control patients converted from negative to positive and two KS patients and one control patient reverted from positive to negative between visits. The remaining 7 KS patients and 20 control patients were negative at both visits.

For the 5 KS patients that converted from an initial negative PBMC result to a positive result at or near to KS diagnosis, the median length of time between the first sample and the KS diagnosis was 19 months. Three of the 6 KS patients that were negative at both visits had their last PBMC sample drawn 2-3 months prior to onset of illness. It is unknown whether these patients became infected between their last study visit and the KS diagnosis date.

DISCUSSION

Ambroziak and coworkers have found evidence that KSHV preferentially infects CD19+ B cells by PBMC subset examination of three patients [19]. Other gammaherpesviruses, such as Epstein-Barr virus (EBV) and herpesvirus saimiri are also lymphotropic herpesviruses

and can cause lymphoproliferative disorders in primates [11, 20].

It is possible that KSHV, like most human herpesviruses, is a ubiquitous infection of adults [21]. EBV, for example, is detectable by PCR in CD19+ B lymphocytes from virtually all seropositive persons [22] and approximately 98% MACS study participants had EBV VCA antibodies at entry into the cohort study [23]. The findings, however, are most consistent with control patients having lower KSHV infection rates than cases and that KSHV is specifically associated with the subsequent development of KS. While it is possible that control patients are infected but have an undetectably low KSHV viral PBMC load, the inability to find evidence of infection in control patients under a variety of PCR conditions suggests that the majority of control patients are not infected. Nonetheless, approximately 10% of these patients were KSHV infected and did not develop KS. It is unknown whether or not this is similar to the KSHV infection rate for the general human population.

This study demonstrates that KSHV infection is both strongly associated with KS and precedes onset of disease in the majority of patients. 57% of KS patients had detectable KSHV infection at their second follow-up visit (52% prior to the onset of KS) compared to only 9% of homosexual/bisexual ( $p=0.002$ ) and 11% of hemophilic control patients ( $p=0.005$ ). Despite similar CD4+ levels between homosexual/bisexual KS cases and controls, KSHV DNA positivity rates were significantly higher for cases at both the first ( $p=0.005$ ) and second sample visits indicating that immunosuppression alone was not responsible for these elevated detection rates. It is also unlikely that KSHV simply colonizes existing KS lesions in AIDS patients since neither patient group had KS at the time the initial sample was obtained. Five KS patients and two homosexual/bisexual control patients



converted from a negative to a positive, possibly due to new infection acquired during the study period.

5 The findings are in contrast to PCR detection of KSHV DNA in all 10 PBMC samples from KS patients by Ambroziak et al. [19]. It is possible that the assay was not sensitive enough to detect virus in all samples since it was required that each positive sample to be repeatedly positive by two independent primers in blinded PCR assays. This appears unlikely, however, given the sensitivity of the PCR nested primer sets. The 7 KS patients who were persistently negative on both paired samples may represent an aviremic or low viral load subpopulation of KS patients. The PCR conditions test a DNA amount equivalent to approximately  $2 \times 10^3$  lymphocytes; an average viral load less than 1 copy per  $2 \times 10^3$  cells may be negative in the assay. Two KS patients and a homosexual/bisexual control patient initially positive for KSHV PCR amplification reverted to negative in samples drawn after diagnosis. These results probably reflect inability to detect KSHV DNA in peripheral blood rather than true loss of infection although more detailed studies of the natural history of infection are needed.

10  
15  
20  
25  
30 The study was designed to answer the fundamental question of whether or not infection with KSHV precedes development of the KS phenotype. The findings indicate that there is a strong antecedent association between KSHV infection and KS. This temporal relationship is an absolute requirement for establishing that KSHV is central to the causal pathway for developing KS. This study contributes additional evidence for a possible causal role for this virus in the development of KS.

REFERENCES

1. Katz MH, Hessol NA, Buchbinder SP, Hirozawa A, O'Malley P, Holmberg SD. Temporal trends of opportunistic infections and malignancies in homosexual men with AIDS. *J Infect Dis.* 1994;170:198-202.

2. Beral V, Peterman TA, Berkelman RL, Jaffe HW. Kaposi's sarcoma among persons with AIDS: a sexually transmitted infection? *Lancet.* 1990;335:123-128.

3. Archibald CP, Schechter MT, Le TN, Craib KJP, Montaner JSG, O'Shaughnessy MV. Evidence for a sexually transmitted cofactor for AIDS-related Kaposi's sarcoma in a cohort of homosexual men. *Epidemiol.* 1992;3:203-209.

4. Beral V, Bull D, Jaffe H, Evans B, Gill N, Tillett H et al. Is risk of Kaposi's sarcoma in AIDS patients in Britain increased if sexual partners came from United States or Africa? *BMJ.* 1991;302:624-5.

5. Beral V. Epidemiology of Kaposi's sarcoma. *Cancer, HIV and AIDS.* London: Imperial Cancer Research Fund; 1991:5-22.

6. Chang Y, Cesarman E, Pessin MS, Lee F, Culpepper J, Knowles DM, et al. Identification of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. *Science.* 1994;265:1865-69.

7. Moore PS, Chang Y. Detection of herpesvirus-like DNA sequences in Kaposi's sarcoma lesions from persons with and without HIV infection. *New England J Med.* 1995;332:1181-1185.

DDBEAD-BZFZB60

- 5
8. Boshoff C, Whitby D, Hatzionnou T, Fisher C, van der Walt J, Hatzakis A et al. Kaposi's sarcoma-associated herpesvirus in HIV-negative Kaposi's sarcoma. *Lancet*. 1995;345:1043-44.
- 10
9. Su I-J, Hsu Y-S, Chang Y-C, Wang I-W. Herpesvirus-like DNA sequence in Kaposi's sarcoma from AIDS and non-AIDS patients in Taiwan. *Lancet*. 1995;345:722-23.
- 15
10. Dupin N, Grandadam M, Calvez V, Gorin I, Aubin JT, Harvard S, et al. Herpesvirus-like DNA in patients with Mediterranean Kaposi's sarcoma. *Lancet*. 1995;345:761-2.
- 20
11. Miller G. Oncogenicity of Epstein-Barr virus. *J Infect Dis*. 1974;130:187-205.
12. Hill AB. Environment and disease: association or causation? *Proc Roy Soc Med*. 1965;58:295-300.
- 25
13. Susser M. Judgment and causal inference: criteria in epidemiologic studies. *Am J Epid*. 1977;105:1-15.
- 30
14. Fishbein DB, Kaplan JE, Spira TJ, Miller B, Schonberger LB, Pinsky PF, et al. Unexplained lymphadenopathy in homosexual men: a longitudinal study. *JAMA*. 1985;254:930-5.
- 35
15. Holmberg SD. Possible cofactors for the development of AIDS-related neoplasms. *Cancer Detection and Prevention*. 1990;14:331-336.
16. Kaslow RA, Ostrow DG, Detels R, Phair JP, Polk BF, Rinaldo CR. The Multicenter AIDS Cohort Study: rationale, organization and selected characteristics of the participants. *Am J Epidemiol*. 1987;126:310-318.

000000000000000000000000000000000000

- 5 17. Wolinsky S, Rinaldo C, Kwok S, Sinsky J, Gupta P, Imagawa D, et al. Human immunodeficiency virus type 1 (HIV-1) infection a median of 18 months before a diagnostic Western blot. *Ann Internal Med.* 1989;111:961.
- 10 18. Bauer HM, Ting Y, Greer CE, Chambers JC, Tashiro CJ, Chimera J, et al. Genital papillomavirus infection in female university students as determined by a PCR-based method. *JAMA.* 1991;265:2809-10.
- 15 19. Ambroziak JA, Blackbourn DJ, Herndier BG, Glogau RG, Gullett JH, McDonald AR, et al. Herpes-like sequences in HIV-infected and uninfected Kaposi's sarcoma patients. *Science.* 1995;268:582-583.
- 20 20. Roizman B. The family Herpesviridae. In: Roizman B, Whitley RJ, Lopez C, eds. *The Human Herpesviruses.* New York: Raven Press, Ltd.; 1993:1-9.
21. Roizman B. New viral footprints in Kaposi's sarcoma. *N Engl J Med.* 1995;332:1227-1228.
- 25 22. Miyashita EM, Yang B, Lam KMC, Crawford DH, Thorley-Lawson DA. A novel form of Epstein-Barr virus latency in normal B cells in vivo. *Cell.* 1995;80:593-601.
- 30 23. Rinaldo CR, Kingsley LA, Lyter DW, Rabin BS, Atchison RW, Bodner AJ, et al. Association of HTLV-III with Epstein-Barr virus infection and abnormalities of T lymphocytes in homosexual men. *J Infect Dis.* 1986;154:556-61.

000000-0270900

EXPERIMENTAL DETAILS SECTION IV:

To determine if the KHV-KS virus is also present in both endemic and HIV-associated KS lesions from African patients, formalin-fixed, paraffin-embedded tissues from both HIV seropositive and HIV seronegative Ugandan KS patients were compared to cancer tissues from patients without KS in a blinded case-control study.

Patient Enrollment: Archival KS biopsy specimens were selected from approximately equal numbers of HIV-associated and endemic HIV-negative KS patients enrolled in an ongoing case-control study of cancer and HIV infection at Makerere University, Kampala Uganda. Control tissues were consecutive archival biopsies from patients with various malignancies enrolled in the same study, chosen without prior knowledge of HIV serostatus. All patients were tested for HIV antibody (measured by Cambridge Bioscience Recombigen Elisa assay).

Tissue preparation: Each sample examined was from an individual patient. Approximately ten tissue sections were cut (10 micron) from each paraffin block using a cleaned knife blade for each specimen. Tissue sections were deparaffinized by extracting the sections twice with 1 ml xylene for 15 min. followed by two extractions with 100% ethanol for 15 min. The remaining pellet was then resuspended and incubated overnight at 50° C in 0.5 ml of lysis buffer (25 mM KCl, 10 mM Tris-HCl, pH 8.3, 1.4 mM MgCl<sub>2</sub>, 0.01% gelatin, 1 mg/ml proteinase K). DNA was extracted with phenol/chloroform, ethanol precipitated and resuspended in 10 mM Tris-HCl, 0.1 mM EDTA, pH 8.3.

PCR Amplification: 0.2-0.4 ug of DNA was used in PCR reactions with KS330<sub>233</sub> primers as previously described [7]. The samples which were negative were retested by nested PCR amplification, which is approximately 10<sup>2</sup>-10<sup>3</sup> fold more sensitive in detecting KS330<sub>233</sub> sequence than

the previously published KS330<sub>233</sub> primer set [7]. These samples were tested twice and samples showing discordant results were retested a third time. 51 of 74 samples initially examined were available for independent  
5 extraction and testing at Chester Beatty Laboratories, London using identical nested PCR primers and conditions to ensure fidelity of the PCR results. Results from eight samples were discordant between laboratories and were removed from the analysis as uninterpretable (four  
10 positive samples from each laboratory). Statistical comparisons were made using EPI-INFO ver. 5 (USD, Stone Mt. GA, USA) with exact confidence intervals.

#### RESULTS:

15 Of 66 tissues examined, 24 were from AIDS-KS cases, 20 were from endemic HIV seronegative KS cases, and 22 were from cancer control patients without KS. Seven of the cancer control patients were HIV seropositive and 15 were HIV seronegative (Figure 22). Tumors examined in  
20 the control group included carcinomas of the breast, ovaries, rectum, stomach, and colon, fibrosarcoma, lymphocytic lymphomas, Hodgkin's lymphomas, choriocarcinoma and anaplastic carcinoma of unknown primary site. The median age of AIDS-KS patients was 29  
25 years (range 3-50) compared to 36 years (range 3-79) for endemic KS patients and 38 years (range 21-73) for cancer controls.

Among KS lesions, 39 of 44 (89%) were positive for  
30 KS330<sub>233</sub> PCR product, including KS tissues from 22 of 24 (92%) HIV seropositive and 17 of 20 (85%) HIV seronegative patients. In comparison, 3 of 22 (14%) nonKS cancer control tissues were positive, including  
1 of 7 (14%) HIV seropositive and 2 of 15 (13%) HIV  
35 seronegative control patients (Figure 19). These control patients included a 73 year old HIV seronegative male and a 29 year old HIV seronegative female with breast carcinomas, and a 36 year old HIV seropositive female with ovarian carcinoma. The odds ratios for

detecting the sequences in tissues from HIV seropositive and HIV seronegative cases and controls was 66 (95% confidence interval (95% C.I.) 3.8-3161) and 36.8 (95% C.I. 4.3-428) respectively. The overall weighted Mantel-Haenzel odds ratio stratified by HIV serostatus was 49.2 (95% C.I. 9.1-335). KS tissues from four HIV seropositive children (ages 3, 5, 6, and 7 years) and four HIV seronegative children (ages 3, 4, 4, and 12 years) were all positive for KS330<sub>233</sub>.

All discordant results (i.e. KSHV negative KS or KSHV positive nonKS cancers) were reviewed microscopically. All KS330<sub>233</sub> PCR negative KS samples were confirmed to be KS. Likewise, all KS330<sub>233</sub> PCR positive nonKS cancers were found not to have occult KS histopathologically.

#### DISCUSSION

These results indicate that KSHV DNA sequences are found not only in AIDS-KS [5], classical KS [6] and transplant KS [7] but also in African KS from both HIV seropositive and seronegative patients. Despite differences in clinical and epidemiological features, KSHV DNA sequences are present in all major clinical subtypes of KS from widely dispersed geographic settings.

This study was performed on banked, formalin-fixed tissues which prevented the use of specific detection assays such as Southern hybridization. DNA extracted after such treatment is often fragmented which reduces the detection sensitivity of PCR and may account for the 5 PCR negative KS samples found in the study. The results, however, are unlikely to be due to PCR contamination or nonspecific amplification. Specimens were tested blindly and a subset of samples were independently extracted and tested at a physically separate laboratory. Specimen blinding is essential to ensure the integrity of results based solely on PCR analyses. A subset of amplicons was sequenced and found to be more than 98% identical to the published KS330<sub>233</sub>

sequence confirming their specific nature and, because of minor sequence variation, making the possibility of contamination unlikely.

5 In contrast to previous studies in North American and European populations, it was found 3 of 22 control tissues to have evidence of KSHV infection. Since these cancers represent a variety of tissue types, it is unlikely that KSHV has an etiologic role in these tumors. One possible explanation for the findings is that these results reflect the rate of KSHV infection in the nonKS population in Uganda. Four independent controlled studies from North America [5 and 9] Europe [7] and Asia [8] have failed to detect evidence of KSHV infection in over 200 cancer control tissues, with the exception of an unusual AIDS-associated, body-cavity-based lymphoma [9]. Taken together, these studies indicate that DNA-based detection of KSHV infection is rare in most nonKS cancer tissues from developed countries. KSHV infection has been reported in post-transplant skin tumors, although well-controlled studies are needed to confirm that these findings are not due to PCR contamination [10]. Since the rate of HIV-negative KS is much more frequent in Uganda than the United States, detection of KSHV in control tissues from cancer patients in the study may reflect a relatively high prevalence infection in the general Ugandan population.

10  
15  
20  
25

While KS is extremely rare among children in developed countries [2], the rate of KS in Ugandan children has risen dramatically over the past 3 decades: age-standardized rates (per 100,000) for boys age 0-14 years were 0.25 in 1964-68 and 10.1 in 1992-93. Detection of KSHV genome in KS lesions from prepubertal children suggests that the virus has a nonsexual mode of transmission among Ugandan children. That five of these children were 5 years old or less raises the possibility that the agent can be transmitted perinatally. Whether or not immune tolerance due to perinatal transmission

30  
35





9. Cesarman E., Chang Y., Moore P.S., Said J.W., Knowles D.M. Kaposi's sarcoma-associated herpesvirus-like DNA sequences are present in AIDS-related body cavity based lymphomas. New England J Med 1995; 332:1186-1191.
10. Rady P.L., et al. Herpesvirus-like DNA sequences in nonKaposi's sarcoma skin lesions of transplant patients. Lancet 1995;345:1339-40.

#### EXPERIMENTAL DETAILS SECTION V:

##### Serologic marker for KSHV infection.

#### METHODS

Patients Serum was collected from a convenience sample of 89 HIV-infected patients seen at several clinical sites in Connecticut, New York, and California. Demographic and clinical information was recorded on standardized forms which were linked to samples by a numerical code. Patients were classified as having KS if the diagnosis was histologically confirmed or, in the opinion of the primary clinician, the diagnosis of KS was unequivocal on clinical grounds. Eighty six (97%) were male; 90 of the 86 men (93%) were homosexual or bisexual. Forty seven patients, all male, had KS. The characteristics of the study population are found in Figure 23].

Cell lines The BCBL-1 line was established from an AIDS-associated body cavity B cell non-Hodgkin's lymphoma [30]. Neither BCBL-1 cells, nor the tumor from which they were derived, express surface immunoglobulin or B cell specific surface markers; however BCBL-1 cells contain immunoglobulin gene rearrangements that are characteristic of B cells [31]. KSHV DNA sequences can be detected in BCBL-1 cells by DNA representational difference analysis [23,32]. BCBL-1 cells also contain

an EBV genome detectable with several different EBV DNA probes. B95-8 is an EBV producer marmoset cell line that can be efficiently induced into EBV lytic cycle gene expression by phorbol esters (TPA) [33,34]. HH 514-16 is an EBV containing cell line, originally from a Burkitt lymphoma, that is optimally inducible into EBV lytic cycle gene expression by n-butyrate [35,36]. B141 is an EBV-negative Burkitt lymphoma cell line [37]. B95-8, HH514-16 and BL41 do not hybridize with the KSHV probes. All cell lines were cultured in RPMI 1640 medium containing 8% fetal calf serum.

Immunoblotting Assays Extracts of uninduced BCBL-1 cells or BCBL-1 cells that had been treated with 20ng/ml TPA and 3 mM n-butyrate for 48 hrs were prepared by sonication. HH514-16 cells, treated similarly, served to control for antibody reactivity to EBV polypeptides. Each lane of a 10% or 12% polyacrylamide gel was loaded with extract of  $5 \times 10^5$  cells in SDS sample buffer; electrophoresis, transfer to nitrocellulose and blocking with skim milk followed standard protocols [38]. Sera were screened at 1:100 dilution. The reaction was developed by 1.0  $\mu$ Ci of  $^{125}$ I Staphylococcal protein A. Radioautographs were exposed to film for 24-48 hrs. Immunoblotting assays were performed and interpreted on coded sera.

Immunofluorescent assay The antigens were BCBL-1 cells that were untreated or treated with 3mM n-butyrate for 48 hrs. Cells were dropped onto slides that were fixed in acetone and methanol. Sera were tested at 1:10 dilution, followed by 1:30 dilution of fluoresceinated goat anti-human Ig. The reactivity of a serum was compared on untreated and n-butyrate treated BCBL-1 cells. Reactivity with 30-50% of the chemically treated BCBL-1 cells was considered a positive reaction. All immunofluorescence tests were performed on coded sera. The two readers were blinded to disease status or results of immunoblotting assays.



and appeared after treatment with chemicals they likely represented lytic cycle rather than latent cycle polypeptides of KSHV.

5     p40 and p60 are KSHV specific: Figures 27A-27B shows  
that antigenic polypeptides corresponding in molecular  
weight to p40 were not observed in two EBV producer  
lines, B95-8 and HH514-16, that were induced into the  
10     EBV lytic cycle by the same chemicals or in comparably  
treated EBV-negative BL41 cells. Furthermore n-butyrate  
strongly induced expression of p40 in BCBL-1 cells but  
had little or no effect on the level of expression of  
the EBV p21 complex in the same cells. In related  
15     experiments it was found that n-butyrate also induced an  
increase in the abundance of KSHV DNA and KSHV lytic  
cycle mRNA. TPA, by contrast, induced the EBV lytic  
cycle efficiently' treatment with TPA caused an increase  
in the abundance of the EBV p21 protein and minimal  
20     induction of KSHV p40. These findings suggested that  
latency to lytic cycle switch of the two gamma herpes  
viruses carried by BCBL-1 cells was under separate  
control and that the p40 complex was specific to the  
KSHV genome.

25     p40 as a serologic marker for KSHV: While a few highly  
reactive sera, such as KS 01-03, (Figure 27B) recognized  
multiple antigenic proteins unique to the chemically  
induced BCBL-1 cells, including p27, p60 and p40, sera  
30     from other patients with KS did not react with p27 or  
p60 but still recognized p40 (Figure 28A and 28B).  
Therefore recognition of p40 was investigated as a  
serologic marker for infection with KSHV. Sera from 89  
HIV-1 infected patients from Connecticut, New York and  
California were examined for presence of antibodies to  
35     p40; only 3 of 42 patients (7%) without KS had  
antibodies to p40 ( $p < 0.0001$  by Chi square). These three  
patients were homosexual or bisexual men from New York  
city. The positive and negative predictive values of  
the serologic marker for the presence of KS were 84% and

78% respectively. Three HIV-1 infected men from New York with non-Hodgkin's lymphoma but without KS were non-reactive to the KSHV p40 antigen. Figure 25 compares the patients with KS whose serum did or did not contain antibodies to KSHV p40. Neither CD4 cell number nor the extent of KS disease predicted the presence or absence of a serologic response to p40.

Immunofluorescence assays: Immunoblots showed that n-butyrate induced expression of KSHV lytic cycle polypeptides in BCBL-1 cells without significantly affecting expression of EBV polypeptides (Figure 28A). Therefore it was reasoned that n-butyrate might also induce many more BCBL-1 cells into the KSHV lytic cycle than into the EBV lytic cycle. Using indirect immunofluorescence with a reference human antiserum, RM in Figure 27B, that contains antibodies to EBV but not KSHV there were about 2% antigen positive untreated BCBL-1 cells and a similar number of antigen positive BCBL cell that had been treated with n-butyrate. Serum 01-03 that is EBV-positive and KS-positive (Figure 27B) detected 2% antigen positive cells in the untreated BCBL population, presumably the EBV expressing cells, while it detected 50% antigen positive BCBL-1 cells that had been treated with n-butyrate. This increase in the number of antigen positive BCBL-1 cells among the n-butyrate treated population served as the basis of an immunofluorescence screening assay for antibodies to KSHV lytic cycle antigens (Figures 29A-29F). The results of the immunofluorescence assay were nearly identical to the immunoblotting assay (Figure 26). Among 89 sera there were only 4 (3%) that were discordant in the two assays. Three sera scored positive by IFA and negative by immunoblotting: one was considered positive by immunoblotting and negative by IFA. 68% of patients with KS and 12% of HIV-1 infected patients without KS were reactive by indirect immunofluorescence assay (IFA). Thus using two different assays, antibodies to KSHV lytic cycle

antigens were found 6 to 9 times more frequently among patients with KS than among HIV-1 infected patients without KS. Stated another way, among individuals who were seropositive to KSHV p40 32/35 (91%) had KS. Among those seropositive by the immunofluorescence assay 32/37 (86%) had KS. Thus infection with KSHV, as defined by these serologic markers, carries a high risk of development of KS.

10 **DISCUSSION**

The recent discovery of genetic sequences representative of a new human herpes virus in KS tumor tissue, taken together with past epidemiologic observations, strongly implicate this novel agent in the pathogenesis of KS. However, these observations, by themselves, do not permit the construction of a unified theory of pathogenesis that accounts for the many mysterious features of KS. For example, the relative contribution of HIV-1, other forms of immunosuppression, geographic factors, sex differences, the role of cytokines and growth factors, and the occurrence of distinct clinical variants must all be eventually understood. By identifying the infection rate in different populations a serologic marker for infection with KSHV would be great aid in unravelling the significance of the new virus in this complicated puzzle.

One possibility is that KSHV, the putative etiologic agent is, like all the other human herpes viruses, a ubiquitous, or at least widespread virus which infects large segments of the human population. Individuals who are immunosuppressed would have a greater likelihood of developing disease, whereas immunocompetent individuals would remain healthy. This pathogenetic model is similar to that postulated for the role that EBV plays in non-Hodgkin's lymphoma or cytomegalovirus in retinitis in patients with AIDS. If this model is correct a very high proportion of the adult human

population might be found to be seropositive for KSHV. The model of a ubiquitous virus selectively causing disease in immunodeficient individuals does not account for classical KS affecting patients who are not immunocompromised nor does it account for the observations that endemic KS in Africa preceded the HIV-1 epidemic. Since many African patients with KS are HIV-1 negative other co-factors must be implicated.

The other possibility is that KSHV infection occurs selectively in the human population. Transmission may be promoted by sexual behavior that also carries a high risk of acquiring HIV-1. In this scenario seroprevalence of KSHV would be expected to be higher in HIV-1 seropositive and HIV-1 seronegative homosexual men than in other populations. If the virus alone were capable of inducing disease, acquisition of KSHV infection, as monitored by the presence of antibody, would be associated with a high rate of clinically evident KS. However, if KSHV infection needed to be accompanied by other co-factors to cause disease, the prevalence of antibody of KSHV might be similar among patients with and without KS. The other co-factors would not be identified in a serologic test for antibodies to KSHV antigens.

The findings, using tests for antibodies to KSHV lytic cycle antigens, are consistent with the general model in which infection with KSHV is infrequent but associated with a high rate of apparent disease. Only a few HIV-1 infected patients without KS had antibodies to the KSHV lytic cycle antigens; by contrast a very high proportion of HIV-1 infected men who had clinically evident KS were seropositive. This finding suggests that a high proportion of individuals who are dually infected with HIV-1 and KSHV develop KS. However, another interpretation of the data is possible, though this interpretation is novel and no other examples are known among the human herpes virus family. Infection with



KSHV might be ubiquitous, antibodies to the virus would not normally be detected in healthy infected individuals. Antibodies would only appear after the virus has been reactivated from the latent into the lytic cycle as might occur during the course of immunosuppression. Thus the two serologic tests that are described would indicate reactivated infection but would not be an index of past exposure to the virus. If this interpretation is correct, it should be possible to demonstrate KSHV DNA sequences or to isolate the virus from healthy individuals who are KSHV seronegative.

Regardless of which of these two interpretations is correct, the serologic studies provide a strong correlation between the presence of antibodies to KSHV lytic cycle gene products and clinical KS. Nonetheless there are two groups of patients whose serologic results require further explanation. One group consists of the few patients with positive serology for KSHV p40 without clinical KS. They may have subclinical or visceral disease, or they may develop KS in the future. The other group is the approximately 30% of patients with KS whose sera lacked antibody to p40. The patients with KS who were p40 seronegative were not misclassified since the diagnosis was confirmed in all of them by biopsy (Figure 25). It is possible that the antibodies being measured are variable and wax and wane with time following infection. The appearance of antibody to p40 may reflect the extent of lytic viral replication which may vary during different phases of the disease. To determine whether this is true prospective studies including serial bleedings are required.

p40 is likely to be only one among a number of KSHV antigens recognized by the infected patients. Antibody recognition of other KSHV antigens may not be possible on immunoblots because they comigrate with EBV polypeptides, because the BCBL-1 cells cannot be induced to express these antigens, or because the antigens are

of low abundance or denatured on the immunoblots. In some individuals serum antibodies to p40 may be consumed in immune complexes with p40 antigen in the circulation. Thus detection of p40 on immunoblots may not be of optimal sensitivity. In this connection three sera recognized antigens in immunofluorescence tests but did not react with p40 on western blots. The serologic test employing whole BCBL-1 cells as antigen are clearly first generation assays to be improved by better characterization of the KSHV gene products and preparation of recombinant antigens.

Lack of a serologic response to p40 could also reflect severely impaired humoral immunity. Although humoral immunity is usually relatively intact in HIV infection, examples of impaired antibody response have been described. For instance, some individuals are known to have impaired antibody responses to parvovirus B19(40 and others have been observed to lose antibodies to hepatitis B surface antigen [41]. An association between the degree of immunosuppression, as monitored by the number of CD4 cells, and the presence or absence of antibody p40 among patients with KS was not found (Figure 25). Furthermore all the patients with or without antibodies to KSHV p40 had antibodies to EBV p21 suggesting an intact humoral immune response.

In these serologic studies, as in the genetic probe studies previously reported, KSHV infection was found in the majority, but not all, patients with KS. Assuming that methodologic explanations do not account exclusively for the seronegative patients, other pathways, in addition to infection with KSHV, may lead to development of KS. In fact, most data suggest that the pathogenesis of KS is a multifactorial process. It has been observed that the product of the HIV-tat gene stimulates growth of KS tissue culture cells [42] and can induce KS-like lesions in mice [43]. These findings suggest a direct role for HIV-1 in the pathogenesis of

5 KS, at least in HIV-infected hosts. In other settings,  
other growth factors may play a similar or complementary  
function. Interleukin-6 and basic fibroblast growth  
factor are both known stimulate growth of KS cells  
10 invitro [44]. Interleukin-6 is also produced in AIDS-KS  
derived cell culture [44]. Thus, KS pathogenesis may  
involve autocrine and paracrine growth factors together  
with infection with KSHV in some patients or with  
certain strains of HIV-1 in other patients. If  
15 infection with KSHV is the *sine qua non* of this process  
on would expect to see evidence of KSHV infection in all  
patients with KS.

20 In summary, an immunoblotting and a immunofluorescence  
screening assay for detection of antibodies to lytic  
cycle antigens of KSHV is disclosed. These assays  
should permit detailed seroepidemiologic investigations  
of KSHV. The findings support the notion of a strong  
association between infection with KSHV and the  
development of KS in HIV-infected patients. Infection  
with KSHV, as defined by these serologic assays, appears  
to carry an extremely high risk of development of  
clinical KS.

006250\* 6474990

REFERENCES:

- 5
1. Kaposi M. Idiopathic multiple pigmented sarcoma of the skin. Cancer 1892; 31:3
2. Safai B, Good RA. Kaposi's sarcoma: A review and recent developments. Clin Bull 1980; 10:62-8.
- 10
3. Loethe R. Kaposi's sarcoma in Ugandan Africans. Acta Pathol Microbiol Scand 1963 (suppl); 161:1-70.
4. Gordon JA. Clinical features of Kaposi's sarcoma amongst Rhodesian Africans. Central African Journal of Medicine 1972; 19:1-6.
- 15
5. Penn I. Kaposi's sarcoma in organ transplant recipients. Transplantation 1979; 27:8-1.
- 20
6. Friedman-Kien AE. Kaposi's sarcoma and Pneumocystis carinii pneumonia among homosexual men - New York City and California. MMWR. 1981; 30:305-8.
- 25
7. Friedman-Kien AE, Laubensin LJ, Rubenstein P, et al. Disseminated Kaposi's sarcoma in homosexual men. Ann Intern Med 1982; 96:693-700.
- 30
8. Berel V, Peterman TA, Berkel RL, Jaffe HW. Kaposi's sarcoma among persons with AIDS: a sexually transmitted disease? Lancet 1990; 335:123-128.
- 35
9. Drew WL, Mills J, Hauer LB, et al. Declining prevalence of Kaposi's sarcoma in homosexual AIDS patients paralleled by a fall in cytomegalovirus transmission. Lancet 1988; 66.
10. Beral V, Peterman TA, Berkelman RL, et al. Kaposi's sarcoma among persons with AIDS: A sexually transmitted infection? Lancet 1990; 2:123.

- 5
11. Barley AC, Downng RG, Cheingson-Popov R et al. HTLV-III serology distinguishes atypical and endemic Kaposi's sarcoma in Africa. *Lancet* 1985; 1:359.
- 10
12. Ziegler, J. L., J.A. Beckstead, P.A. Volberding, et al. 1984. Non-Hodgkin's lymphoma in 90 homosexual men: Relation to generalized lymphadenopathy and the acquired immunodeficiency syndrome. *N. Engl. J. Med.* 311:565-570.
- 15
13. Levine AM. Non-Hodgkins lymphomas and other malifnancies in the acquired immunodeficiency syndrome. *Semin Oncol* 1987; 14:34-9.
- 20
14. Knowles DM, Chamulak M, Subar M, et al. Lymphoid neoplasia associated with the acquired immunodeficiency syndrome. *Ann Intern Med* 1988; 108:744-53.
- 25
15. MacMahon EME, Glass JD, Hayward SD, Mann RB, Becher PS, Charache P, McArthur JC, Ambender RF. Epstein-Barr virus in AIDS-related primary central nervous system lymphoma. *Lancet* 1991; 338:969-73.
- 30
16. Sillman FH, Sedlis A. Anogenital papillomavirus infection adn neoplasia in immunodeficiency women. *Obstet Gynecol Clin North Am* 1987; 260:348-53.
- 35
17. Drew WL, Mills J. Hauer LB, Miner RC, et al. Cytomegalovirus and Kaposi's sarcoma in young homosexual men. *Lancet.* 1982; 2:125-127.
18. Callant JE, Moore RD, Richman DD, Keruly J, Chaisson RE. Risk factors for Kaposi's sarcoma in patients with advanced human immunodeficiency virus disease treated wiht zidovudine. *Arch Inter Med* 1994; 154:566-72.

03290702950

- 5
- 10
- 15
- 20
- 25
- 30
- 35
19. Giraldo, G., E. Beth, and E. Huang. 1980. Kaposi's sarcoma and its relationship to cytomegalovirus III CMV DNA and CMV early antigens in Kaposi's sarcoma. *Int. J. Cancer* 26:23-29.
  20. Ambinder, R.F., Newman, C., Hawyard, G.S. et al. 1987 Lack of association of cytomegalovirus with endemic African Kaposi's sarcoma *J. Inf. Dis.* 156:193-7.
  21. Jahan N, Razzaque A, Greenspan J et al. Analysis of human KS biopsies and cloned cell lines for cytomegalovirus, HIV-1 and other selected DNA virus sequences. *ADIS Res Human Retrovirus* 1989; 5:225.
  22. Wang RY-H, Shih, JW-k, Weiss SH, et al. Mycoplasma penetrans infections in male homosexuals with AIDS: high seroprevalence and association with Kaposi's sarcoma. *Clin Infect Dis* 1993; 17:724-29.
  23. Chang YE, Cesarman E, Pessin MS, Lee F, Culpepper J, Knowles DM, Moore PS. Identification of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. *Science* 1994: 266; 1865-9.
  24. Moore PS, Cahng Y. Dectection of Herpes virus-like DNA sequences in Kaposi's sarcoma inpateints with and those without HIV infection. *N Engl J Med* 1995; 332"1181-5.
  25. Su IJ, Hsu YS, Chang YC, Wang IW. Herpesvirus-like DNA sequences in Kaposi's sarcoma from AIDS and non-AIDS pateints in Taiwan. *Lancet* 1995; 345:722-3.
  26. Huang YQ, Li JJ, Kaplan MH, Polesz B, Katabira WC, Zhang D, et al. Human herpesvirus-like nucleic acid in various forms of Kaposi's sarcoma. *Lancet* 1995; 345:759-61.

27. Dupin N, Grandadam M, Calvez V, Gorin I, Aubin JT, Harvard S, et al. Herpesvirus-like DNA sequence in patients with Mediterranean Kaposi's sarcoma. Lancet 1995; 345:761-2.
- 5
28. Collandre H, Ferris S, Grau O, Montagnier L, Blanchard A, Kaposi's sarcoma and new herpes virus. Lancet 1995; 345:1043.
- 10
29. Boshoff C, Whitby D, Hatzioannu T, Fisher C, van der Walt J, Hatzakis A, Weiss R, Schultz T. Kaposi's sarcoma - associated herpes virus in HIV-negative Kaposi's Sarcoma. Lancet 1995; 345:1043-4.
- 15
30. Cesarman E, Chang Y, Moore PS, Said JW, Knowles DM. Kaposi's sarcoma associated herpes virus-like DNA sequences in AIDS-related body-cavity-based lymphomas. N Engl J Med 1995; 332:1186-91.
- 20
31. Knowles, D. M., G. Inghirami, A. Ubriaco, and R. Dalla-Favera. 1989. Molecular genetic analysis of three AIDS-associated neoplasms of uncertain lineage demonstrate their B-cell derivation and the possible pathogenetic role of the Epstein-Barr virus. Blood 74:792-799.
- 25
32. Lisitsyn, N., N. Lisitsyn, and M. Wigler. 1993. Cloning the differences between two complex genomes. Science 259:946-951.
- 30
33. Miller G, Lipman M, Release of infectious Epstein-Barr virus by transformed marmoset leukocytes. Proc Nat Acad Sci USA 1973; 70:190-4.
- 35
34. zur Hausen, H., F. J. O'Neill, U. K. Freese, and E. Hecker. 1978. Persisting oncogenic herpesvirus induced by tumor promoter TPA. Naturer 272:373-375.

35. Rabson, M., L. Heston, and G, Miller. 1983. Identification of a rare Epstein-Barr virus variant which enhances early antigen expression in Raji cells. Proc. Natl. aCad. Sci. USA 80:2762-2766.
- 5
36. Luka, J., B. Kallin, and G. Klien. 1979. Induction of the Epstein-Barr virus (EBV) cycle in latently infected cells by n-butyrate. Virology 94:228-231.
- 10
37. Calendar A, Billaud M, Aubry J-P, Bauchereau J, Vuillaume M, Lenoir GM. Epstein-Barr virus (EBV) induces expression of B-cell activation markers on in vitro infection of EBV-neative B-lymphoma cells. Proc nat Acad Sci USA 1987; 84;8060-4.
- 15
38. Twobin, H., Staehelin, T., Gordon, J. Electrophoeretic transfer of proteins from polyacrylamide gels to nitrocullulose sheets: procedure and some applications Proc Natl Acad Sci USA 1979; 76: 4350-4354.
- 20
39. van Grunsven, W.M.J., E. C. van Heerde, H.J.W. de Haard, W.J.M. Spaan, and J Middledorp. 1993. Gene mapping and expression of two immunodominant Epstein-Barr virus capsid proteins. J. Virol. 67:3908-3916.
- 25
40. Frickhofen N, Abkowitz JL, Safford M, Berry M, Antunez-de-Mayolo J, et al. Persistent B19 Parvovirus infection in pateints infected with human immunodeficiency virus type 1 (HIV-1): a treatable cause of anemia in ADIS. Ann Intern Med 1990; 113:926-33.
- 30
41. Biggar RJ, Goedert JJ, Hoofnagle J: Accelerated loss of antibody to hepatitis B surface antigen among immunodeficient homosexual men infected with HIV. N Engl J Med 1987; 316:630-31.
- 35

006290" 62 FZ 950



42. Ensoli B, Barillari G, Salahuddin SZ et al. Tat protein of HIV-1 stimulates growth of cells derived from Kaposi's sarcoma lesions of AIDS patients. Nature 1990; 345:84.
- 5
43. Vogel J, Hinrichs SH, Reynolds RK, Luciw PA, Jay G. The HIV tat gene induces dermal lesions resembling Kaposi's sarcoma in transgenic mice. Nature 1994; 335:601-11.
- 10
44. Miles SA, Rezai AR, Salazar-Gonzalez JF, Stevens RH, Logan DM, Mistuyasu RT, Taga T, et al. AIDS Kaposi's sarcoma-derived cells produce and respond to interleukin-6. Proc Natl Acad Sci USA 1990; 87: 4068-72.
- 15
- 006290\* 62 F 20 96 8